



Laboratory tests for diagnosing and monitoring canine leishmaniasis

Paltrinieri, S ; Gradoni, L ; Roura, X ; Zatelli, A ; Zini, Eric

Abstract: Although several reviews on canine leishmaniasis have been published, none thoroughly described clinicopathologic abnormalities and their clinical usefulness. The aim of this review was to provide information concerning current diagnostic tests relevant for clinical pathologists and from a practical perspective. Specifically, in canine leishmaniasis, nonregenerative normocytic normochromic anemia, thrombocytopenia, or leukogram changes may be present. Clinical chemistry and urinalysis may indicate renal dysfunction (azotemia, decreased urine specific gravity, proteinuria) and an inflammatory/immune response (increased acute phase proteins [APP] or 2 - and/or -globulins). Although a potential gammopathy is usually polyclonal, it may also appear oligo- or monoclonal, especially in dogs coinfecting by other vector-borne pathogens. When lesions are accessible to fine-needle aspiration (lymphadenomegaly, nodular lesions, joint swelling), cytology is strongly advised, as the presence of *Leishmania* amastigotes in a pattern of pyogranulomatous inflammation or lymphoplasmacytic hyperplasia is diagnostic. If the cytologic pattern is inconclusive, the parasite should be identified by histology/immunohistochemistry or PCR on surgical biopsies. Alternatively, cytology and PCR may be performed on bone marrow samples where amastigotes, along with erythroid hypoplasia, myeloid hyperplasia, plasmacytosis, or secondary dysmyelopoiesis can be observed. Dogs with overt leishmaniasis generally have high antibody titers, while low titers predominate in immunologically resistant infected dogs or in exposed dogs with no parasite confirmation. Quantitative serology is recommended in clinically suspect dogs as high-titer antibodies titers may confirm the clinical diagnosis. In confirmed and treated dogs, renal function and inflammatory/immune response variables should be periodically monitored.

DOI: <https://doi.org/10.1111/vcp.12413>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-128082>

Journal Article

Accepted Version

Originally published at:

Paltrinieri, S; Gradoni, L; Roura, X; Zatelli, A; Zini, Eric (2016). Laboratory tests for diagnosing and monitoring canine leishmaniasis. *Veterinary Clinical Pathology*, 45(4):552-578.

DOI: <https://doi.org/10.1111/vcp.12413>

1 ***LABORATORY TESTS FOR DIAGNOSING AND MONITORING CANINE LEISHMANIASIS***

2

3

4 **Running header: laboratory diagnosis of leishmaniasis**

5

6 Saverio Paltrinieri, DVM, PhD, Dipl ECVCP, Department of Veterinary Sciences and Public

7 Health – University of Milan, Italy

8 Luigi Gradoni, Unit of Vector-borne Diseases & International Health, Department of Infectious

9 Diseases, Istituto Superiore di Sanità, Rome, Italy

10 Xavier Roura, Hospital Clínic Veterinari, Universitat Autònoma de Barcelona, Bellaterra, Spain;

11 Andrea Zatelli, Medical Consultancy Services, TàXbiex, Malta

12 Eric Zini, Clinic for Small Animal Internal Medicine, University of Zurich, Switzerland;

13 Department of Animal Medicine, Production and Health, University of Padova, Italy; Istituto

14 Veterinario di Novara, Granozzo con Monticello (NO), Italy.

15

16 Corresponding author:

17 Prof. Saverio Paltrinieri

18 Dipartimento di Scienze Veterinarie e Sanità Pubblica, Università di Milano

19 Via Celoria 10, 20133, Milano, Italy

20 Tel 0039-02-50318103

21 Fax 0039-02-50318095

22 e-mail: saverio.paltrinieri@unimi.it

23

24 **Abstract**

25 Despite several reviews on canine leishmaniasis, none throughly described laboratory abnormalities
26 and diagnostic tests in light of their clinical usefulness. Aim was to summarize this information
27 from a practical perspective. Canine leishmaniasis often leads to profound clinicopathological
28 abnormalities. CBC reveals non-regenerative normocytic normochromic anemia and
29 thrombocytopenia, but other types of anemia or leukogram changes are also observed. Clinical
30 chemistry and urinalysis frequently show abnormalities suggestive of renal dysfunction (increased
31 creatinine and urea, decreased urine specific gravity, proteinuria) and inflammatory/immune
32 activation (increased acute phase proteins or α_2 - and/or γ -globulins). The gammopathy is typically
33 polyclonal but can also be oligo- or monoclonal. When lesions accessible to fine needle aspiration
34 or other sampling techniques (lymphadenomegaly, nodular or ulcerated skin lesions, joint
35 swelling) are present, cytology is strongly advised; the disease is confirmed if *Leishmania*
36 amastigotes are identified and pyogranulmatous inflammation or lymphoplasmocytic hyperplasia is
37 present. If results are inconclusive, the parasite should be identified through
38 histology/immunohistochemistry or PCR on surgical biopsies. Alternatively, cytology and PCR
39 may be performed on bone marrow, where amastigotes and erythroid hypoplasia, myeloid
40 hyperplasia, plasmocytosis, or secondary dysmyelopoiesis can be observed. Dogs with overt disease
41 generally have high antibody titers at ELISA or immunofluorescence. Low antibody titers may be
42 found in infected dogs that are immunologically resistant, or in exposed dogs (dogs not harboring
43 the parasite). If leishmaniasis is suspected but cytological evidence of the parasite is negative,
44 serology is recommended for confirmation. In treated dogs, renal function and the magnitude of the
45 inflammatory/immune reaction should be periodically monitored.

46

47

48 ***Keywords: Dog; Leishmania infantum; clinical usefulness; diagnosis; follow-up***

49

51 Leishmaniasis is a frequent infectious disease in dogs living in endemic areas, is associated with
52 important morbidity and, although appropriate treatment, it can cause death. Despite several
53 reviews have been published so far, none has fully described the diagnostic role of laboratory
54 findings that are observed in dogs with leishmaniasis and of available tests. Therefore, the aim of
55 the present review was to provide information concerning laboratory abnormalities and current
56 diagnostic tests, from a practical perspective.

57

58 ***Etiology and pathogenesis of canine leishmaniasis***

59 Canine leishmaniasis is a disease caused by the protozoan parasite *Leishmania infantum*
60 (Kinetoplastida: Trypanosomatidae), or its New World synonym *Leishmania chagasi*.¹ Although
61 non-vectorial transmission is occasionally reported (e.g. by transplacental, transfusional or venereal
62 route),²⁻⁴ it is assumed that canine leishmaniasis is mainly acquired by the bite of infected
63 phlebotomine sand flies (Diptera: Psychodidae). Therefore, the geographical distribution and
64 prevalence of the disease largely depends on the presence and abundance of competent vector
65 species, including members of the *Phlebotomus* (*Larroussius*) subgenus in the Old World
66 (Mediterranean Basin and Central Asia) and *Lutzomyia longipalpis* and *L. evansi* in the New World
67 (from Mexico through Argentina).⁵ Blood-sucking females ingest the intracellular, non-flagellated
68 form of *Leishmania* (amastigote) during the blood meal on infected hosts. After multiplication,
69 flagellated forms (promastigote) and transform into non-replicative, infectious metacyclic forms
70 that are inoculated into the host's dermis when the fly takes another blood meal. Parasites are
71 phagocytosed by local and recruited host cells, and within phagolysosomes of resident macrophages
72 promastigotes transform and replicate as amastigotes.⁶

73 The pathogenesis of the disease starts with the ingestion of recently inoculated promastigotes by
74 neutrophils that are unable to kill the protozoa but activate dermal macrophages and other
75 mechanisms of innate immunity (e.g. acute phase response). Macrophages initially activate their
76 antimicrobial defenses, which are based on the release of enzymes and on the production of free

radicals within the phagolysosome. *Leishmania* organisms are able to interfere with the oxidative activity of the phagocytes^{7,8} and therefore amastigotes can survive and actively replicate in macrophages, leading to cell destruction and infecting progressively more and more phagocytes. In longitudinal field studies employing naïve dogs, it has been demonstrated that *Leishmania* can be detected by PCR in bone marrow samples starting about 6 months from natural exposure to phlebotomine vectors.⁹ Once bone marrow has been colonized it is generally accepted that the dog is persistently infected. However, the same studies showed that dogs being PCR positive in bone marrow may become negative in the following months without any treatment; it is unknown whether the parasite density fluctuates below the threshold limit of the test, the infection persists in organs other than bone marrow, or the host defenses successfully eradicate the infection.⁹ Despite dogs can mount an antibody response after the first contact with the parasites, resistance or susceptibility to progressive infection (leading to clinical disease) depend on the balance between Th1 (cell-mediated) and Th2 (humoral) immune responses of the host: those with prevailing Th2 response are more likely to have uncontrolled dissemination of parasites to all body tissues and to show lesions and clinical signs of leishmaniasis.¹⁰⁻¹³

Of note, depending on the host-parasite interactions, the simple detection of circulating antibodies does not necessarily imply that the dog is actually infected and detection of parasites in tissues does not suggest that the infected dog is actually sick. Therefore, the two existing guidelines that deal with the diagnosis and staging of canine leishmaniasis, recently released by the Leishvet group¹⁴ and the Canine Leishmaniasis Working Group (CLWG),¹⁵ classify dogs as exposed, infected or sick based on a combination of clinical findings and clinicopathological tests or tests for etiological diagnosis. Specifically, both guidelines include two main categories of dogs:

- Infected dogs: dogs clinically unremarkable, without abnormalities in hematology, clinical chemistry and urinalysis, that test positive to PCR or cytology in tissues such as bone marrow, lymph node, spleen, skin or peripheral blood;

- Sick dogs: infected dogs that have clinical signs or clinicopathological changes consistent with leishmaniasis.

The classification of the CLWG¹⁵ includes two additional categories of dogs at the extremes of the spectrum:

- Exposed dogs: dogs clinically unremarkable with positive serology, in which PCR or cytology fail to demonstrate the presence of the parasite;
- Severely sick dogs: sick dogs that do not respond to treatments and/or are affected by concurrent diseases.

The Leishvet group classifies sick dogs in four stages according to the severity of clinical signs and magnitude of antileishmanial antibody titers.¹⁴

Clinical signs of canine leishmaniasis

The interpretation of clinicopathological, serological and molecular tests should be done in light of history, signalment and clinical presentation. History should focus on the risk of exposure to phlebotomine vectors. Signalment should focus on gender, breed and age; clinically overt leishmaniasis is more frequent in male dogs, Boxer and German shepherd, and in those younger than 3 years or older than 8 years,^{16,17} although recent reports suggests that any dog above 2 years is at risk.¹⁸

The spectrum of clinical presentations is wide and ranges from infections characterized by the absence of obvious clinical findings but detectable laboratory abnormalities, to overt clinical infections characterized by the presence of clinical and laboratory abnormalities that require or not hospitalization. However, some signs are more frequent than others. The most typical clinical findings include:^{14,15,19-22} generalized lymphadenomegaly, cutaneous lesions, pale mucous membranes, weight loss or cachexia, polyuria/polydipsia, epistaxis, onicogriphosis. Ocular lesions,^{15,23,24} lameness, lethargy and fever are also quite common.^{25,26} In a longitudinal study,²⁷ it was shown that lymph node enlargement appears first (approximately 1 year after exposure to sand

flies), followed by cutaneous lesions (after 18 months) and ocular signs (after 22 months). Furthermore, atypical forms have been described, especially in highly endemic areas, including gastrointestinal, neurological, musculoskeletal, cardiopulmonary, lower urinary tract or genital tract signs.²⁸⁻³⁶

Laboratory abnormalities that may support or confirm leishmaniasis

In addition to clinical findings, laboratory abnormalities detectable by routine hematology, clinical chemistry or urinalysis may further increase the clinical suspicion of canine leishmaniasis. Moreover, especially in the early phases of the disease or with atypical presentations, laboratory changes may occur in the absence of obvious abnormalities at physical examination. Therefore, a basic panel is mandatory when canine leishmaniasis is suspected or when a dog with positive result tests for etiological diagnosis needs to be classified as “exposed”, “infected” or “sick”. Table 1 summarizes the clinicopathological changes that are found in dogs with leishmaniasis (i.e. “sick” dogs).

1) Hematologic abnormalities

Hematological changes in canine leishmaniasis are non specific.^{37,38} Neutrophilia, due to the systemic inflammatory response associated with the infectious diseases, may be particularly prominent if ulcerative cutaneous lesions with secondary bacterial infection are present.^{37,39,40} Conversely, numerical or morphological changes in the other leukocyte populations are less common, although lymphopenia, lymphocytosis or eosinophilia are occasionally described³⁹⁻⁴² Amastigotes may be rarely documented in circulating leukocytes of infected dogs (0.5% of cases), within neutrophils, lymphocytes and monocytes (figure 1);^{41,43} the percentage of infected cells is so low that their search is generally not rewarding. When a systemic disease and blood dissemination is suspected, more sensitive tests such as PCR or quantitative PCR should be preferred to the simple microscopical examination of blood smears (see below). When parasites are found on smears,

154 however, it is very likely that the parasite burden is high: therefore, although, as stated above, the
155 simple detection of the parasite does not automatically mean that the infected dog is sick, it is more
156 likely that dogs with a high parasite burden (and parasites detected on smears) are sick than simply
157 infected.

158 The most common hematological changes in leishmaniotic dogs is anemia,^{37,38,44,45} that may appear
159 6 months after exposure to infection.²⁷ Anemia is usually mild to moderate and has the normocytic
160 normochromic non regenerative pattern^{40,44,45} typical of the anemia of inflammatory disease, whose
161 pathogenesis include a depression of metabolic activity of bone marrow, in this case sustained also
162 by the infection of stromal cells by the parasite, and a decreased iron availability for
163 erythropoiesis.⁴⁶ However the pathogenesis of anemia in leishmaniotic dogs include additional
164 mechanisms such renal failure leading to reduced erythropoietin synthesis and secondary
165 dismyelopoiesis associated with the production of antibodies against erythroid precursors.⁴⁷
166 Moreover, it is very likely that anemia also has a hemolytic component,⁴⁸ likely due to the presence
167 of anti-RBC antibodies; in a few cases, the hemolysis may be prevalent,⁴⁹ or associated with a
168 “lupus-like” reaction along with other dermatological or systemic signs or laboratory change,
169 including positive ANA-test⁵⁰ or perinuclear antineutrophil cytoplasmic autoantibodies.⁵¹

170 Thrombocytopenia is fairly frequent in leishmaniotic dogs without concurrent infections. It is
171 usually mild to moderate but if severe, co-infections with other vector-borne pathogens (e.g.
172 *Ehrlichia canis*, *Anaplasma phagocytophilum* or *A. platys*) or other possible causes of reduced
173 platelet concentration should be suspected. The most likely mechanism responsible for
174 thrombocytopenia in leishmaniasis is a peripheral consumption of circulating platelets. In turn, this
175 mechanism may be due to immune-mediated mechanisms, since the presence of anti-Plt antibodies
176 has been demonstrated in leishmaniotic dogs.⁵²⁻⁵⁴

177 Moreover, platelet loss may be associated to hypercoagulability caused by a decreased
178 concentration of anti-thrombin III, as in any other protein losing nephropathy⁵⁵ (see below), or to
179 disseminated intravascular coagulation (DIC) that may be occasionally suspected in leishmaniotic

180 dogs.⁵⁶ However, the mechanism of thrombocytopenia in leishmaniotic dogs includes also a
181 decreased production due to the depressed bone marrow activity and to the secondary
182 dysmyelopoiesis cited above. Even in the absence of reduced platelet counts, however, platelet may
183 be hypofunctional in dogs with leishmaniasis⁵⁷ although this reduced function is rarely responsible
184 for hemostatic abnormalities, as described below.

185 One additional hematological test that may be run in dogs with leishmaniasis is the flow-cytometric
186 evaluation of the CD4/CD8 ratio. The rationale for performing this test is that as soon as Th1
187 responses decreases, thus increasing the susceptibility to the disease and favoring the shift from
188 latent infection to overt disease, the number of CD4+ lymphocytes decreases causing reduction of
189 the CD4/CD8 ratio.^{58,59} Therefore, a seropositive or PCR-positive dog with a low CD4/CD8 ratio is
190 more predisposed to develop clinical signs than a similar dog with normal CD4/CD8 ratio. The
191 practical applicability of this test, however, is limited by the high individual variability and by the
192 difficulty to determine a clear cut-off for staging the disease. Hence, this test may be useful for
193 monitoring the post-treatment follow-up, rather than to stage a dog at first diagnosis of
194 leishmaniasis; the authors do not suggest the use of this test for diagnostic purposes in dogs
195 suspected to have leishmaniasis.

196 Finally, the hematological profile of leishmaniotic dogs may be completed by bone marrow
197 cytology.^{27,42,44,60} This analysis may be useful in leishmaniotic dogs to confirm the infection
198 through the detection of infected macrophages, as better specified below, but it may be also used to
199 differentiate a simple infection from systemic disease (i.e. “infected” vs. “sick” dog).¹⁵ Despite
200 some histological studies demonstrated that parasite density can be high despite few clinical signs,⁶¹
201 generally the parasite load and the presence of cytological alterations increase as soon as the dogs
202 show clinical signs.⁶² Therefore, rare infected macrophages may be occasionally seen in the absence
203 of other pathological findings in dogs that are simply infected, whereas dogs in which the parasite
204 induces an inflammatory reaction and/or a systemic disease, are characterized by a higher number
205 of parasites detected cytologically and by a series of morphological changes. Specifically, in the

206 latter case cytology of the bone marrow usually reveals an erythroid hypoplasia,⁴⁰ without
207 abnormalities in the ratio between maturative and proliferative pools of erythroid precursors,
208 occasionally associated with myeloid hyperplasia (and thus with an increased M:E ratio). Moreover,
209 signs of bone marrow inflammation, generically defined by Stockham and Scott as “myelitis”,⁶³, are
210 usually found (figure 2). These include a proliferation of either infected or non-infected
211 macrophages often with signs of erythrophagia or, more generically, cytophagia, associated with an
212 increase of neutrophils in the framework of the myeloid hyperplasia, and a moderate to severe
213 plasmocytosis characterized by a higher number of plasma cells with signs of activation (e.g. mott
214 cells), and lymphocytes.^{64,65} Megakaryocyte hyperplasia may also be present, especially when
215 peripheral consumption of platelets occurs.

216 Secondary dysmyelopoiesis is another common finding, although less frequent than the typical
217 pattern listed above (figure 3). This condition is characterized by multiple peripheral cytopenias
218 (e.g. the anemia and thrombocytopenia cited above) associated with hypercellular bone marrow on
219 which one or more cell lineages show dysplastic features. In canine leishmaniasis, these mostly
220 include dyserythropoiesis and dysmegakariopoiesis, while dysmyelopoiesis (and in particular
221 dysgranulopoiesis, mostly characterized by abnormal maturation and ring forms) is only
222 occasionally found: more specifically, erythroid precursors with abnormal mitoses, asynchronous
223 nucleo-cytoplasmic maturation, nuclear fragmentation, and/or late stage maturation arrest,
224 especially when the pathogenesis includes an immune-mediated reaction against RBC precursors.

225 Dwarf megakaryocytes associated or not with an increased number of megakarioblasts and with
226 emperipolesis may be typical aspects of dysmegakariopoiesis in leishmaniotic dogs. The detection
227 of secondary dysmyelopoiesis however, is not per se diagnostic for leishmaniasis, unless
228 macrophages with intracytoplasmic amastigotes and other signs of reactivity (e.g.
229 lymphoplasmocytic infiltration) are found. Therefore, the cause-effect association between
230 secondary dysmyelopoiesis and the simple seropositivity or PCR-positivity should be carefully
231 considered. Ultimately, in this case the diagnosis of leishmaniasis should be based on the exclusion

232 of other infectious, toxic or metabolic causes of secondary dysmyelopoiesis or of primary
233 myelodysplastic syndromes (MDS) as recommended in hematology textbooks.^{45,47,64}
234 The last and less common finding in bone marrow cytology leishmaniotic dogs may be the presence
235 of erythroid hyperplasia when peripheral signs are consistent with the diagnosis of immune-
236 mediated hemolytic anemia (IMHA). Also in this case, however, IMHA can be associated with
237 leishmaniasis only if the parasite is found in cytological specimens characterized by erythroid
238 hyperplasia. If not, the diagnostic approach should be taken into account only if any other possible
239 cause of IMHA has been excluded. In brief, bone marrow cytology may be useful for diagnostic
240 purposes in some dogs, by detecting amastigotes and compatible cytological abnormalities, or to
241 differentiate between infected dogs from those that are sick due to leishmaniasis.

242

243 2) *Hemostatic abnormalities*

244 Hemostatic abnormalities are uncommon in leishmaniotic dogs. Activated partial thromboplastin
245 time (aPTT) and prothrombin time (PT) may be increased; however, in most cases this is due to
246 preanalytical factors, since their prolongation may occur when the concentration of total globulins
247 increases, which is frequent in dogs with leishmaniasis.⁶⁶ Alternatively, prolonged coagulation
248 times may result from DIC, although this complication is uncommon in leishmaniotic dogs.^{56,67}
249 Conversely, hypercoagulability may be common in leishmaniotic dogs if affected by severe protein
250 losing nephropathy.⁶⁸ This is mostly due to glomerular loss of antithrombin III (ATIII), a protease
251 inhibitor involved in the regulation of blood coagulation that prevents the conversion of fibrinogen
252 into fibrin. The lack of this physiologic anticoagulant may induce hypercoagulability that in turn
253 promotes thrombosis and subsequent consumption coagulopathy. Hypercoagulability is also
254 favored by the hyperviscosity syndrome due to increased circulating globulins.⁶⁹ As in any other
255 hypercoagulable state associated with protein losing nephropathy, hypercoagulability of
256 leishmaniotic dogs was also demonstrated through a decreased clot formation time and an increased
257 global clot strength using thromboelastography (TEG);⁷⁰ differently, in another study the

258 coagulation profile of leishmaniotic dogs assessed by thromboelastometry (TEM, a technique
259 similar to TEG) was within normal limits.⁷¹ However, it is worth noting that TEM and TEG are
260 affected by the RBC mass, possibly explaining the different results obtained with TEM and
261 TEG.^{72,73} In brief, to assess hypercoagulability in dogs with protein losing nephropathy associated
262 with leishmaniasis the authors currently suggest including only ATIII measurement.

263

264 3) *Biochemical abnormalities*

265 Because the clinical presentation of dogs with leishmaniasis is variable, also the type of
266 biochemical abnormalities varies accordingly. Renal dysfunction and inflammation/immune
267 reactions are frequently observed and their presence should be evaluated in each dog with suspected
268 or confirmed leishmaniasis (see below). Regarding other routine biochemical analytes that may be
269 altered in leishmaniotic dogs, there are enzymes of hepatobiliary or pancreatic damage that may
270 increase in case of pyogranulomatous infiltrates affecting these organs,^{15,16} muscular enzymes,
271 including LDH and CK, that may increase with musculoskeletal lesion.⁷⁴ Nevertheless, increased
272 CK may also be due to the increased CK-BB isoenzyme if neurological signs are present,⁷⁵ since
273 *Leishmania* has been found in the brain of some affected dogs with cerebrovascular alterations,⁷⁶⁻⁷⁸
274 or to increased CK-MB if leishmaniotic dogs have cardiomyopathy (i.e. increased troponin I and
275 cardiopulmonary lesions have been reported in many leishmaniotic dogs).⁷⁹⁻⁸¹ Biochemical
276 abnormalities consistent with alteration of endocrine organs are rare, despite amastigotes associated
277 with inflammatory lesions have been found in the adrenal cortex of leishmaniotic dogs.^{82,83}

278

279 *Assessment of renal function*

280 The systemic immune complex disease that characterizes leishmaniasis induces the deposition of
281 circulating immune complexes at the glomerular level. This induces a series of inflammatory
282 glomerular changes detectable histologically and ultramicroscopically,^{36,84,85,86,87} that are primarily
283 responsible for a proteinuric nephropathy.⁸⁷ The evolution of this condition, as for any other

protein-losing nephropathy, is represented by the development of a chronic kidney disease (CKD) due to a series of factors that include progressive glomerulosclerosis, renal hypertension, overload of protein reabsorption in tubular cells, with subsequent tubulointerstitial nephritis^{85,87} In turn, advanced stages of CKD are characterized by hyperazotemia and may be associated with systemic hypertension, both factors contributing to comorbidity in dogs with leishmaniasis.^{87,88} Therefore, as for any other renal disease, the early detection of proteinuria and correct classification of CKD is mandatory in the diagnostic workup. To this aim, the clinical and laboratory approach is the same recommended by the International Renal Interest Society (IRIS)⁸⁹ for any type of CKD. This approach is based on a thorough clinical evaluation of the dog, based on physical examination and diagnostic imaging, on the measurement of arterial pressure according to the guidelines recommended by the American College of Veterinary Internal Medicine (ACVIM)⁹⁰ and on the quantification of urinary proteins (described in the section of this article regarding urinalysis) and markers of renal function. According to the IRIS guidelines, these include mostly the urine specific gravity and the serum concentration of creatinine.⁸⁹ This latter has been shown to increase frequently in leishmaniotic dogs^{14,15,16,19,20,91} and early (from 12 to 18 months) following infection.²⁷ Creatinine is a good indicator of a decreased glomerular filtration rate (GFR), and it more specific than other markers such as urea^{92,93} but unfortunately they are not enough sensitive to detect the earliest stages of renal insufficiency. Due to the compensatory activity of residual nephrons, in fact, an increase of serum creatinine occur only when the GFR dramatically decreases.⁹³ Therefore, although creatinine is commonly used to detect overt CKD, a huge research activity is currently running in veterinary nephrology and clinical pathology to identify earlier markers of decreased GFR, either in leishmaniotic dogs or in dogs affected by other types of CKD. The direct measurement of GFR trough clearance tests (e.g. clearance of inulin, exogenous creatinine or, more recently, iohexol) would be the best method to assess in real time the functionality of the kidneys^{94,95} Unfortunately these tests, although currently validated for use in dogs, are not widely available in clinical practices or in veterinary laboratories. Therefore, the diagnostic potential of

indirect markers other than creatinine has been recently investigated. Cystatin C (Cys C) has been proposed as surrogate marker of creatinine, especially using advanced measurement techniques such as particle-enhanced turbidimetric immunoassay.⁹⁶⁻⁹⁹ The serum concentration of Cys C has been assessed also in dogs with leishmaniasis.¹⁰⁰ However, there is no evidence that serum Cys C, that is less specific than creatinine, is more sensitive than creatinine in detecting early CKD.⁹⁴ Promising results have been obtained by the measurement of urinary Cys C¹⁰¹ but this seems not to be true in canine leishmaniasis, where urinary Cys C increases only in proteinuric dogs with increased creatinine.¹⁰² Recently, a promising biomarker for early detection of renal dysfunction seems the symmetric dimethylarginine (SDMA), a catabolite of methylated proteins that is mainly excreted by the glomeruli and not reabsorbed by tubuli,^{103,104} that in published report has been measured in serum using chromatographic techniques.¹⁰⁴ Studies in cats, where the early diagnosis of CKD may be more challenging than in dogs, demonstrated that SDMA increases earlier than creatinine,^{105,106} although it does not provide additional advantages in animals with overt CKD (i.e. with CKD on which serum creatinine is already increased). Preliminary results indicate that SDMA may have the same role of early indicator of renal dysfunction also in dogs,^{104,107} although a few extrarenal variables may affect the results.¹⁰⁸ Therefore, although no studies specifically focused on the use of SDMA in canine leishmaniasis exist, it is very likely that SDMA may be used, in the future, to assess renal function in leishmaniotic dogs that are proteinuric but still have normal creatinine concentration.

Other blood markers may provide additional information in leishmaniotic patients with CKD. For example in people the increased serum concentration of homocysteine (Hcy), endothelin-1 (ET-1) or C-reactive protein (CRP) may predict, hypertension and/or inflammation, especially in association with CKD.¹⁰⁹⁻¹¹³ In dogs, Hcy increases in those with CKD¹¹⁴ but seems to be less associated with hypertension than in humans.¹¹⁵ Conversely, ET-1 increases in serum of dogs with hypertension associated with early stages of CKD¹¹⁵ and is also associated with inflammation, as demonstrated by its correlation with CRP levels, which in turn increases mostly in CKD associated

336 with inflammatory conditions.¹¹³ However, the potential of these additional markers as
337 complementary tests for the management of leishmaniotic dogs with CKD has not been
338 investigated. Conversely, inflammatory markers such as CRP, ferritin and adiponectin have been
339 measured in the urine of dogs with leishmaniasis and were found increased in dogs with proteinuria,
340 sometime in the absence of elevated serum creatinine.^{102,116,117} These analytes might therefore work
341 as early markers of CKD in leishmaniotic dogs, although it is more likely that their increase depend
342 on the elevated serum concentration, thus reflecting the inflammatory state typical of leishmaniasis,
343 rather than a direct consequence of CKD.

344 Finally, in leishmaniotic dogs, tubulointerstitial lesions may occur secondarily to proteinuria caused
345 by glomerular damages. Therefore, the availability of markers to identify dogs with tubular lesions
346 would allow to early differentiate dogs with pure glomerular lesions from dogs with more advanced
347 renal damage. Tubular markers, however, are usually measured in urine and not in serum and are
348 described in the section on urinalysis. It is also worth mentioning that some dogs with CKD may
349 have acute deterioration of their renal dysfunction due to factors related or not to leishmaniasis
350 (e.g. vomiting, diarrhea).

351

352 *Assessment of inflammatory/immune reactions*

353 Based on the pathophysiology above described, it is clear that leishmaniotic dogs with overt disease
354 have an intense inflammatory reaction and produces high amount of molecules involved in the
355 immune response, including antibodies. Both these phenomena may be investigated using tests such
356 as protein analysis and serum protein electrophoresis or measurement of acute phase proteins
357 (APPs).

358

359 Protein analysis and serum protein electrophoresis are considered the most accurate tests to
360 diagnose canine leishmaniasis. Protein changes appear 12-18 months after exposure to the
361 parasite.²⁷ Total proteins and total globulin are frequently increased, especially in the acute phase of

the disease.^{14,15,16,21,91,118,119} The increase of total protein can correlate with the severity of the clinical score, according to Proverbio et al.¹²⁰ Moreover albumin decreases both because of its role as negative APPs (see below) and of the renal loss associated with proteinuric nephropathy leading to decreased albumin:globulin (A/G) ratio^{91,118,119} The decrease of the A/G ratio is so frequent that it has been considered by some Authors to be one of the more sensitive tests for canine leishmaniasis¹¹⁸ and hypoalbuminemia is considered a negative prognostic factor in leishmaniotic dogs since the magnitude of hypoalbuminemia negatively correlates with survival times.¹²¹ The typical electrophoretogram of leishmaniotic dogs with overt clinical signs (figure 4) displays hypoalbuminemia, an increase of α_2 -globulin, where most of the positive APPs migrate, and especially a strong increase of γ -globulins, due to the huge amount of circulating antibodies, immune complexes, and other molecules with γ motility involved in immune-mediated inflammatory reactions.¹¹⁹ Occasionally, peaks due to circulating antibodies are found in the β region, where IgM and some APPs migrates. The gammopathy is typically polyclonal but sometime the peak may be narrower (oligoclonal), biclonal¹²² or definitely monoclonal.¹²³ The interpretation of the peak must take into account the electrophoretic method, since peaks are usually narrower using capillary zonal electrophoresis (CZE) than using agarose gel or cellulose acetate.¹²⁴ Therefore, CZE may lead to falsely interpret as monoclonal a peak that is ultimately oligo- or polyclonal (figure 5). Moreover, although monoclonal peaks associated exclusively with leishmaniasis have been described, the detection of monoclonal peaks should induce to consider the possible presence of concurrent diseases (e.g. other vector-borne diseases or multiple myeloma).^{125,126}

Acute phase proteins are powerful indicators of inflammation: the pro-inflammatory cytokines produced in inflammatory sites induce the so called “acute phase response”, characterized by the release of neutrophils from storage pools, by an activation of myelopoiesis (see above), and by a

modulation of protein synthesis in the liver.¹¹⁹ This latter phenomenon leads to a decreased serum concentration of the “negative APPs”, and to an increased concentration of the “positive APPs” that includes a series of immunomodulators, scavenger or transport proteins, antiproteases, and other proteins involved in host defenses.¹¹² Therefore it is not surprising that the serum concentration of positive APPs in dogs with overt canine leishmaniasis is high. The list of APPs whose concentration increases in serum of leishmaniotic dogs is long and includes CRP, Haptoglobin (Hp), Ceruloplasmin (Cp) Serum Amyloid A (SAA) and ferritin,¹²⁷⁻¹³² with CRP and ferritin that, as previously reported, increase also in urine. Similarly, a decrease of negative APPs other than albumin has also been reported; these are transferrin, indirectly measured as total iron binding capacity (TIBC), that induces also a reduction in the concentration of iron, and a decreased activity of the enzyme paraoxonase (PON-1).^{129,130,133} PON-1 is a negative APP that is bound to high density lipoproteins (HDL) and represents a link between inflammation and oxidative stress. Therefore its decrease is not constantly seen in leishmaniotic dogs but it may become evident when oxidative stress is particularly severe.¹³³ Interestingly, in these cases also the concentration of HDL, that is converted into low density lipoprotein (LDL) after detachment of PON-1, decreases¹³⁴ and may be a cheap marker of inflammation and oxidative stress associated with leishmaniasis. Recently a reduced serum activity of adenosine deaminase (ADA) and butyrylcholinesterase (BChE), two enzymes involved in modulating immune responses, has also been reported in dogs with leishmaniasis.¹³⁵

The APP changes summarized above are not diagnostic *per se* since mild increases of positive APPs have been reported also in infected dogs without clinical signs¹²⁷ and severely increased levels may occur in diseases other than leishmaniasis.¹¹² In a dog in which leishmaniasis has been diagnosed by other clinical or laboratory findings, however, the magnitude of these changes may reflect the magnitude of inflammation and thus provide prognostic information. In particular, the decrease of PON-1 is particularly evident in severe diseases and may therefore be a negative prognostic marker. Moreover, APPs may be useful to monitor the follow up.

413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438

4) *Abnormalities at urinalysis*

The analysis of urine from dogs with suspected or confirmed leishmaniasis must be focused on three main aspects: confirmation of CKD, evaluation of proteinuria, and investigation of possible tubular damage. Urine samples should be analyzed using the usual approach followed in any case of CKD, that starts with the physico-chemical analysis and continues with sediment analysis and evaluation of proteinuria, eventually followed by the measurement of markers of tubular damages.

Physico-chemical analysis

According to standardized procedures,¹³⁶ after centrifugation of urine to separate supernatants from the sediment, the routine physicochemical analysis must be addressed to evaluate with a refractometer the urine specific gravity (USG), that tends to decrease in dogs with tubulo-interstitial damage.⁹³ Then, the supernatant can be used for a dipstick analysis, that should be focused on assessing:

- the pH that, although not extremely accurate¹³⁷ may be useful to properly interpret other dipstick results: for example the semiquantification of proteins may provide erroneously positive results in alkaline urine (pH>8);⁹³
- the concentration of proteins, to be interpreted as described below;
- the presence of glucose, that, as specified below, may be an indicator of tubular damage.

Sediment analysis is another important step of urinalysis of leishmaniotic dogs: the presence of an active sediment (e.g. a sediment with high numbers of leukocytes, erythrocytes or bacteria) is important either because may indicate a lower urinary tract infection superimposed on the primary disease (leishmaniasis) or because in the presence of an active sediment, results regarding proteinuria should be carefully considered;¹³⁸ conversely, the presence of granular or cellular casts may be consistent with tubular damage.⁹³

439 *Evaluation of proteinuria*

440 The evaluation of proteinuria is an essential step, since the presence and magnitude of proteinuria is
441 not only a marker of kidney disease, but also a risk factor for the progression of nephropathy.¹³⁹
442 According to the ACVIM guidelines,¹⁴⁰ proteinuria should be assessed in any dog with diseases
443 potentially inducing this condition, such as leishmaniasis. The ACVIM guidelines recommend to
444 collect urines by cystocentesis, to avoid contamination from the lower urinary tract, however, to
445 improve the possibility to frequently monitor dogs at risk to develop proteinuria, a first evaluation
446 may be done on voided samples, since results recorded with the two methods of collection overlap
447 when the sediment is inactive.¹⁴¹ Additionally, it may be appropriate to screen the possible presence
448 of proteinuria using a dipstick, since it has been demonstrated that if the dipstick is negative or the
449 dogs is likely non proteinuric according to the IRIS classification⁸⁹ and any additional evaluation of
450 proteinuria is therefore not necessary.¹⁴² Conversely, if the dipstick is positive it is very likely that
451 the dog is proteinuric and the protein to creatinine (UPC) ratio must be run to quantify the level of
452 proteinuria. Similarly the UPC ratio must be measured in dogs with a weakly positive dipstick
453 result but with a low USG, in order to understand if the weak positive result really reflect the
454 presence of proteinuria or not.¹⁴² In all the cases above, the UPC ratio must be measured to
455 definitely classify the dog as proteinuric (UPC >0.5), borderline proteinuric (UPC= 0.2-0.5) or non
456 proteinuric (UPC <0.2) according to the IRIS classification, recently revised for the diagnosis of
457 glomerular disease.^{89,143} In the interpretation of data, particular attention should be paid on results
458 close to these threshold since results may be affected by analytical variability or by the type of
459 reagent or laboratory procedures used to measure urinary proteins.¹⁴⁴⁻¹⁴⁶ Then, especially if the dogs
460 is borderline proteinuric or proteinuric, quantification of proteinuria must be repeatedly assessed (at
461 least 3 times in 2 weeks) because, according to the ACVIM guidelines,¹⁴⁰ additional investigations
462 or treatments should be performed only if persistent proteinuria is confirmed by repeated urinalysis.
463 Alternatively, the analysis of proteinuria can be done on pooled urine samples, in order to reduce
464 the possible influence of circadian variations in renal protein excretion.¹⁴⁷ Finally, according to the

465 ACVIM guidelines¹⁴⁰ the origin of urinary protein should be assessed through a renal biopsy.
466 However, according to the recent IRIS guidelines¹⁴³ renal biopsy is recommended only in the case
467 of rapid progression of CKD or in dogs not responding to conventional treatments. Alternatively,
468 the origin of proteinuria can be argued on the basis of surrogate methods such as qualitative analysis
469 of urinary proteins (see below). However, in the case of very high UPC ratio, as usually occur in
470 leishmaniotic dogs, proteinuria is mostly of glomerular origin and tubular proteinuria, if present,
471 contributes only partially to the concentration of urinary proteins.

472

473 *Markers of tubular injury*

474 Although the gold standard method to identify the origin of proteinuria is the histological analysis
475 of renal biopsies, due to its invasiveness this technique is rarely applied in leishmaniotic dogs.
476 However, in order to differentiate the dogs with a tubular component of proteinuria, that are in a
477 more advanced stage of renal disease, urinary markers may be used.¹⁴⁸ Some rough markers such as
478 granular or cellular casts and glycosuria in normoglycemic dogs are very specific indicators of
479 tubular damage, but are not enough sensitive and therefore do not detect dogs with early tubular
480 damage. Additionally, tubular damage so severe to induce glycosuria are rarely observed in
481 leishmaniotic dogs. Early information about the presence of tubular damage may be achieved using
482 sodium dodecylsulphate (SDS) electrophoresis of urinary proteins or measuring the concentration
483 or some urinary analyte or the activity of some urinary enzymes. The SDS denaturates and charges
484 negatively the urinary proteins. Therefore, after migration on polyacrylamide gel (SDS-PAGE) or
485 agarose gel (SDS-AGE), proteins migrate according their molecular mass.¹⁴⁹ This allows to
486 differentiate large proteins, that are of glomerular origin, from small proteins, that are of tubular
487 origin. Results of SDS-PAGE have been shown to well correlate with results of renal biopsies,
488 especially for the identification of glomerular damage or of severe tubule-interstitial damages.¹⁵⁰⁻¹⁵¹
489 while SDS-AGE, that is more diffuse in veterinary laboratories because it may be fully automated
490 and is more rapid, cheap and less toxic than SDS-PAGE.¹⁵¹ However, SDS-AGE has some

analytical limitations:¹⁵² it is more susceptible to the concentration of urine thus potentially providing false positive or false negative results in very concentrated or very diluted urine, respectively, and may suffer from pre-analytical artifacts such as bands due to storage artifacts or to proteins originating in the male genital tract.^{153,154} However, using SDS-AGE it has been demonstrated that most of the leishmaniotic dogs have a mixed (glomerular and tubular) pattern and only a minority of dogs, likely those with less “chronic” CKD, have a pure glomerular proteinuria.^{84,155} Occasionally, some leishmaniotic dogs has low molecular weight proteinuria with no signs of glomerular disease, likely due to a free light chain proteinuria (i.e. a pre-renal proteinuria associated with the intense antibody production) rather than to a tubular damage.¹⁵⁶ Enzymuria is considered a good marker of tubular damage, since the enzymes of interest are usually located in the cytoplasm of tubular cells. Therefore, they may be found in urine when tubular cells are damaged. Several enzymes may be used for this purpose but the two most popular are γ -glutamyl transferase (GGT) and N-acetyl- β -N-glucosaminidase (NAG)^{93,157} that must be measured just after sampling since their activity decreases after refrigeration or freezing.¹⁵⁸ Increases of these enzymes, as well as of other enzyme such as alkaline phosphatase (ALP) or β -glucuronidase, have been reported in dog with leishmaniasis¹⁵⁹ and the increase of GGT activity has been showed to correlate with the presence of tubular bands in SDS.¹⁵⁵ On the contrary, no information is available on the utility in leishmaniotic dogs of the measurement of other urinary analytes such as clusterin, kidney injury molecule 1 (Kim-1), neutrophil gelatinase-associated lipocalin (NGAL) or retinol binding protein (RBP), used in pharmacology to assess the presence of renal tubular damages and that have been shown to be potentially useful markers in dogs with CKD.^{148,160-162}

512

513 ***Tests for etiological diagnosis that may support or confirm leishmaniasis***

514 Tests for etiological diagnosis are used to identify the presence of the parasite or its components
515 (direct tests) or the host’s response to the parasite (indirect tests). As previously mentioned, positive

indirect tests (serology) may or may not indicate a current infection. Conversely, positive direct tests (cytology, histology, immunohistochemistry, PCR, culture and xenodiagnosis) demonstrate that the dog is actually harboring *Leishmania* and it is therefore infected. However, as stated above, the relationship between infection and disease should be based on the evaluation of clinical findings and clinicopathologic tests. The most common tests for etiological diagnosis are described below.

521

1) Serology

Methods

Apart from some techniques such as Western blotting, that is highly accurate but not available in routine practice, or other methods that have been proposed but are not extensively used, such as latex agglutination or methods based on the detection of antibodies through immunosensors or flow cytometry,¹⁶³⁻¹⁶⁶ the most common techniques used to detect antibodies are based on three analytical principles: immunofluorescent antibody test (IFAT), enzyme-linked immunosorbent assay (ELISA) and immunochromatographic test (ICT). ICT is the basis of all-rapid “in clinic” assays, which have a major limitation being that they provide results in a qualitative manner (i.e. presence/absence of specific reactive bands).¹⁶⁷ Several commercial ICT kits are available, which employ single or multiple recombinant *Leishmania* antigens to be used on serum, plasma, whole blood or blood spots dried onto filter paper.¹⁶⁸ Several studies have evaluated their diagnostic performances and, in general, they reach similar conclusions: while specificity is quite acceptable, ICT sensitivity is usually low (in the approximate range of 30-70%).¹⁶⁹ Lowest sensitivities are found in infected dogs without clinical signs, the highest ones in those with overt disease.¹⁷⁰ Therefore, ICT may be used as a first “in clinic” test to complete the laboratory evaluation of clinically suspected dogs and, in case of positivity, serology should be repeated by ELISA or IFAT, which provide quantitative results (see below). However, due to the low sensitivity of the ICT test, a negative result may be falsely negative and therefore, if the clinical suspicion persists, tests with higher sensitivity (i.e. IFAT or ELISA) should be performed. Recently, a ICT kit claiming detection

of antibodies developed after natural infection but not those elicited by vaccination with the LiESP-based vaccine, has been proposed as a tool to differentiate vaccinated from infected dogs.¹⁷¹ The principle of the test is sound, and the first studies reported an elevated sensitivity of this ICT format;¹⁷² however, other studies reported a low sensitivity also for this test.¹⁷³ IFAT is recognized as the reference method to perform anti-*Leishmania* serology in dogs,^{169,174} as it is very sensitive and also highly specific except in areas endemic for the New World parasite *Trypanosoma cruzi*, that may give false positive results; values approach 100% for both the parameters. ELISA is also very sensitive and specific when a combination of immunodominant, recombinant proteins are used as antigen, whereas it has slightly lower specificity when crude parasite lysates are employed instead.^{167,173-175} Compared to IFAT, that is based on the evaluation of promastigote fluorescence at UV microscope and is therefore operator-dependent), ELISA is easier to standardize since results are read by an automated spectrophotometer. Both IFAT and ELISA have the advantage to provide quantitative results that are based, respectively, on the final antibody titer (the last sample dilution providing positive result) and optical density values that can be compared with reference titred samples using conversion factors. Based on the unavoidable variability due to operator-dependent or analytical (antigen stability, antiserum or equipment performances) parameters, precise anti-*Leishmania* antibody titers are not universally available. Hence, a titer is considered “high” if it is 4 fold higher than the threshold value of the laboratory.¹⁵ Similarly, 4 fold variations in titers of sequential samples of the same dog should be expected in seroconversions, or in the outcome of therapy. Hence, sequential samples must be analyzed by the same method in the same laboratory.

563

564 *Interpretation*

565 Serological tests detect and quantify the presence of antibodies in serum or plasma. Antibodies can
566 be found in blood as soon as 1 month after exposure to infected phlebotomines, although the
567 median time for seroconversion is about 5 months in natural conditions and 3 in experimental

studies).¹⁷⁶ Therefore is very likely that dogs living in highly endemic regions seroconvert during the sand fly activity period (from late spring to early autumn in temperate zones, all over the year in tropical ones),⁹ If the infection is efficiently controlled by the host's immune responses, the antibody titers, when present, tend to remain low and therefore these clinically-healthy dogs are classified as exposed or infected.¹⁵ Conversely, the uncontrolled parasite dissemination is associated with an exaggerated humoral response and therefore antibody titers are high when the disease is evident. This condition is classified as "sick dog" or "severely sick dog" by CLWG classification, and stage III or IV (severe or very severe disease) by Leishvet classification.¹⁴ Furthermore, a direct relationship between the clinical score and antibody titers exists.^{120,177} However, dogs with clinical signs (sick) and low-medium antibody titers may also be detected. These have been classified as stage I or II (mild or moderate disease) according to the Leishvet classification.¹⁴ Therefore, quantitative serology should always be performed when, despite strong clinical suspicion of leishmaniasis, lesions approachable by fine needle aspiration are not present or when cytological analysis of lesions (including lymphoid organs and bone marrow) does not reveal the typical pattern associated with leishmaniasis, despite a possible PCR positivity. In this case a high antibody titer is often consistent with the disease, while, if the antibody titer is low, leishmaniasis should be considered only if any other disease potentially responsible of the clinical presentation is ruled out.^{14,15}

2) PCR

Methods

As for serology, several molecular methods have been proposed to detect the presence of the parasite DNA in various biological samples. Some of these methods are not commonly used or recently validated, such as the non-amplification assay based on the use of probes labelled with gold nanoparticles¹⁷⁸ or the loop-mediated isothermal amplification (LAMP).^{179,180} Conversely, conventional PCR, nested PCR and quantitative (real time) PCR are widely used in routine

594 practice.^{14,15,169,174} PCR sensitivity and specificity varies according to the method and especially
595 according to the target DNA sequence. Most of the PCR tests currently used are targeting multicopy
596 DNA sequences, such as the small subunit ribosomal RNA genes or the kinetoplast DNA
597 minicircles, thus increasing the sensitivity of the test.¹⁸¹ Compared with conventional and nested
598 PCR, the quantitative PCR techniques offer two main advantages:¹⁸² they are usually run in close
599 systems and are therefore less prone to contamination, and provide information about the copies of
600 DNA that are present in the sample. This latter aspect may be relevant during the follow up to
601 monitor the efficacy of leishmanicidal treatments and therefore it may advisable to use quantitative
602 PCR at first diagnosis (before any treatment), in order to have a baseline value for further analyses
603 during the follow up^{182,183} However, it does not seem that quantitative PCR techniques are more
604 sensitive than conventional or nested PCR to diagnose leishmaniasis in dogs.¹⁸⁴ One additional
605 limitation of quantitative PCR is that standardized methods to accurately quantify the copies of
606 DNA may not be offered by some laboratories.

607

608 *Samples*

609 PCR techniques may be applied virtually on any tissue or biological fluids. Theoretically, it may be
610 superfluous to use molecular tests in affected tissues in which amastigotes have been visualized by
611 cytology or histology (see below). However, cytology and histology are less sensitive than PCR
612 (see below) and therefore, a negative cytological result does not exclude that amastigotes are
613 present. Therefore, when a fine needle aspirate or a tissue biopsy is performed, it may be advisable
614 to use part of the material to prepare cytological or histological specimens and to store the
615 remaining sample as recommended by the laboratory (usually frozen, fixed in 95% ethyl alcohol or
616 in appropriate preservatives) to run PCR in case amastigotes cannot be visualized despite the
617 cytological or histological pattern is highly consistent with leishmaniasis. If needed, PCR may also
618 be performed on cytological material already fixed on glass slides¹⁸⁵ or on formalin fixed and
619 paraffin embedded material.^{186,187}

620 In routine practice PCR is rarely run on injured tissues, for which cytology and histology are
621 preferred. When lesions are not present, or they are not approachable by fine needle aspiration or
622 biopsy (for example when the prevalent clinical presentation is anemia or proteinuric nephropathy),
623 other samples are preferred. Several studies demonstrated that bone marrow and/or lymph nodes
624 and spleen provide the highest sensitivity in detecting *Leishmania* by PCR, especially in sick
625 dogs,^{14,188-192} pending that the quality of the sample is adequate, especially for the lymph node.
626 Parasite DNA can be easily found in the skin, including intact tissue especially in the facial region
627 or in the ears. Recent studies demonstrated that conjunctival and, to a lesser extent, oral and nasal
628 swab material is very sensitive for the detection of *Leishmania* DNA and, in addition, can provide
629 positive results earlier than other tissues^{188,190,193-196} Buffy coat or whole blood may also be used for
630 conventional or quantitative PCR analysis. Although its sensitivity is lower than that of the other
631 tissues cited above, it can be collected without invasive methods and in the case of positive results
632 provide a diagnostic information in a rapid and cheap way^{14,15}

633

634 *Interpretation*

635 When interpreting PCR results it must be kept in mind the difference between infected and sick
636 dogs. Ultimately, the detection of the parasite's DNA indicates that the dog is infected. The
637 correlation between infection and disease should be based on the presence of clinical signs and
638 laboratory abnormalities. From this perspective, the detection of *Leishmania* DNA in lesions with
639 cytological or histological patterns highly consistent with leishmaniasis (see below) supports the
640 diagnosis of disease and, similarly, a positive PCR in blood or bone marrow of a dog with systemic
641 signs of leishmaniasis can support the diagnosis of disease. Conversely, positive results in dogs
642 without signs clearly referable to leishmaniasis do not support the hypothesis that the infected dog
643 is also affected by clinical leishmaniasis, unless any other possible disease is excluded. For
644 example, a transient PCR-positivity in bone marrow may be found a few months from the natural
645 exposure to sand fly bites, without necessarily meaning that the dogs is infected or sick.⁹ Similarly,

the PCR positivity in intact skin of a dog living in an endemic area does not necessarily suggest that the presence of parasites in the skin will be followed by parasite dissemination throughout other body tissues.¹⁰⁻¹³ Skin positive PCR results may in fact be dependent on the presence of recently-inoculated promastigotes, or of amastigotes recently phagocytosed by resident macrophages that, in “resistant” dogs, may efficiently control the infection at local level.^{188,192,197}

651

3) Cytology

Samples and methods

Fine needle aspiration should be performed in all cases showing cutaneous papular or nodular lesions and/or lymph node enlargement.¹⁵ Ulcerative cutaneous lesions can be sampled by scraping the lesion or using less invasive methods such as imprint smears. Additionally reports describing the presence of amastigotes and associated lesions in nodular masses with atypical localization, such as the tongue,^{29,33} the testis,¹⁹⁸⁻¹⁹⁹ and oral or nasal masses²⁰⁰ have been reported and therefore any nodular lesion in dogs with clinical or laboratory signs potentially consistent with leishmaniasis (e.g. anemia, CKD, alterations of the electrophoretograms, positive serology) should be sampled by fine needle aspiration. Nasal lesions may also be sampled using brush cytology²⁰¹ Similarly, when clinical or clinicopathological pattern is consistent with leishmaniasis, the possible presence of *Leishmania* should be investigated also in pathological body fluids such as joint fluids,^{25,26} effusions,⁴¹ or cerebrospinal fluid although in this latter sample, cellularity is usually so low that PCR may identify the parasite better than cytology.⁷⁷ In all these cases, fluid samples must be collected by centesis, following the procedures routinely recommended for each site.

When cutaneous lesions or nodular lesions in other organs, lymph node enlargement, abnormal accumulation of fluids are absent but the clinical suspicion of leishmaniasis is high, the possible presence of parasites should be investigated in organs rich of cells of the monocyte-macrophage system, such as bone marrow, lymph nodes or spleen^{14,15,60}

671

672 *Interpretation*

673 Cytology aims to demonstrate the presence of amastigotes that are usually found within the
674 macrophages and, when the parasite burden is high and cell lysis occurs, also on the background
675 (figure 6). The detection of amastigotes may be difficult in cutaneous ulcerative lesions, where the
676 presence of necrosis and cellular debris or of contaminating bacteria may mask the presence of
677 amastigotes. Attention should be paid to misinterpret as amastigotes cellular or granular debris that
678 may be present in these lesions.

679 Additionally, cytology may allow to detect the typical inflammatory patterns associated with
680 leishmaniasis, that are usually characterized by pyogranulomatous inflammation associated with a
681 moderate to severe lymphoplasmocytic infiltration in skin or nodular lesions with atypical
682 localization (figure 5) and, in lymph nodes, by a reactive hyperplasia of variable severity,
683 characterized by lymphoplasmocytic and macrophagic infiltration, usually associated with
684 numerous neutrophils.^{60,202,203} Similarly, cytologic patterns typically associated with leishmaniasis
685 may be found in the bone marrow, as described in details in the section dedicated to hematology.
686 Neutrophils, lymphocytes and macrophages can be found also in body fluids of dogs affected by
687 leishmaniasis. Additionally, the fluids may show the features typical of inflammation such as high
688 cell counts and protein contents and, in joint fluids, decreased viscosity and negative mucin clot
689 test.^{25,26,41,77,204,205}

690 The diagnosis of leishmaniasis is easy when amastigotes are detected in samples that shows the
691 cytologic patterns described above. However, when cytologic patterns consistent with leishmaniasis
692 but no amastigotes are seen, leishmaniasis should not be ruled out, since it is known that the
693 diagnostic sensitivity of cytology is low.^{169,174} In these cases, tests that have higher analytical and
694 diagnostic sensitivity, such as PCR, must be run. Alternatively, affected tissues can be biopsied to
695 perform histology and immunohistochemistry, as described below. Conversely, when amastigotes
696 are seen in the absence of cytological abnormalities, or cytology is done on bone marrow, lymph
697 node or spleen, positive results must be interpreted carefully, as systemic signs may be due to

698 diseases other than leishmaniasis.¹⁵ Similarly, a diagnostic workup to confirm or exclude
699 leishmaniasis (i.e. the clinical disease associated with infection) should be run when *Leishmania* is
700 incidentally found in lesions that clearly have a different origin. For example, several reports
701 describe the association between the presence of amastigotes and tumors such as lymphoma,
702 transmissible venereal tumors and other types of neoplasia.²⁰⁶⁻²¹¹ On a practical standpoint in these
703 cases it is important to understand if the dog is affected by both diseases or affected by a neoplastic
704 disease and simply infected with *Leishmania*.

705

706 4) *Histology*

707 Histology can demonstrate the presence of *Leishmania* in routinely hematoxylin and eosin stained
708 sections when cytology provides parasite-negative results in tissues having a cytological pattern
709 highly consistent with leishmaniasis. Compared with PCR, histology has two main disadvantages: it
710 can be more laborious and time consuming, and the identification of amastigotes may be more
711 difficult than in cytological samples. As for the latter, amastigote presence can be confirmed by
712 immunohistochemistry (figure 7),^{36,212} in situ hybridization^{213,214} or PCR on formalin-fixed and
713 paraffin embedded samples.^{186,187} On the other hand, histology has the advantage to provide
714 additional information on the cytoarchitectural pattern of the lesions. This is a great advantage since
715 it may allow to discriminate dogs in which the parasite is associated with typical lesions from those
716 in which the infection does not seem to be associated with the disease. Therefore, according to some
717 guidelines,²² histology should always be performed, especially in the case of cutaneous lesions. The
718 interpretation of histological results is facilitated by the elevated number of papers describing the
719 distribution of parasites and the lesions associated with active disease, mostly characterized by
720 lymphoplasmacytic or granulomatous-pyogranulomatous inflammations and/or by vasculitis either
721 in organs usually affected by *Leishmania* (bone marrow, spleen, skin, lymph nodes, kidney, etc) but
722 also in unusual sites such as heart, lung, adrenal gland, genital tract, central nervous system, skeletal

723 muscle, gastrointestinal tract, nails, lacrimal glands and ocular muscles.^{23,24,29-31,33,36,74,76-78,80-82,84-}
724 86,156,202,215-221

725

726 5) *Parasite culture and biological test for infectiousness (xenodiagnosis)*

727 Conclusive diagnosis of active infection should be based on tissue cultures, which not only confirm
728 whether dogs harbor parasites, but also demonstrate that the protozoa is viable and multiplies
729 actively. A diagnostic *Leishmania* culture, which differs from maintenance culture of laboratory-
730 adapted strains, requires biphasic blood-agar media that need fresh components.⁷⁵ A conclusive test
731 for infectiousness (xenodiagnosis) requires that naive (laboratory-reared) sand flies are induced to
732 feed on infected dogs and are examined thereafter for the presence of promastigotes in the gut.²²²
733 However both tests are unpractical, since on one hand specific blood-agar media are unavailable
734 commercially for diagnostic *Leishmania* culture and must be prepared in laboratory, on the other
735 hand only a few centers breed sand flies for diagnostic purposes. Therefore these tests are mainly
736 intended for research and cannot be recommended for routine practice.

737

738 ***Future perspectives***

739 Several studies investigated the diagnostic potential of innovative markers in dogs with
740 leishmaniasis: for example, the measurement of iron superoxide dismutase (Fe-SODe) secreted by
741 the parasite has been evaluated as a possible marker of infection;²²³ proteomic analysis revealed a
742 series of proteins that are over- or under-represented in leishmaniotic dogs;²²⁴ the analysis of the
743 expression of cytokines or molecules such as leptin or inducible nitric oxide synthetase (iNOS) in
744 blood or tissues revealed different profiles in leishmaniotic dogs compared to controls^{11,13,225-227}
745 high levels of matrix metalloproteinases (MMP), and especially of MMP9 and pro-MMP2, have
746 been reported in serum or CSF of leishmaniotic dogs.^{228,229} However, none of the studies provided,
747 to date, exhaustive information on the possible utility in practice of these markers. However, the

information provided in these investigations is useful to design future research and explore their potential clinical application.

Similarly, the attention of researchers has been recently focused on markers of oxidative stress; inflammation is characterized by the release of reactive oxygen metabolites from phagocytes recruited in inflammatory sites and this leads to a consumption of antioxidant compounds.²³⁰ Increases of oxidants or oxidized molecules (e.g. reactive oxygen metabolites, malonyldialdehyde, lipoperoxides, thiobarbituric acid reacting substances) and decreases of antioxidant compounds (total antioxidant capacity, trace elements, paraoxonase) have been reported in leishmaniotic dogs^{130,133,134,231-234} Unfortunately, only a few studies assessed the diagnostic potential of oxidative markers. for example, the diagnostic performance of ROS were not excellent, likely because oxidative phenomena are intense when inflammation is superimposed to *Leishmania* infection during the clinically overt phase, but depressed if macrophages have been recently infected, due to the ability of the parasite to inhibit the production of oxygen radicals by phagocytes.^{7,8,234}

Tests for monitoring the post-treatment follow up

Laboratory tests during the follow up should be focused in monitoring possible toxic effect of treatment as well as the clinical and the parasitological status of the patient following administration of drugs according to conventional treatments protocols. These mainly include the administration of antimonials and allopurinol or of miltefosine and allopurinol. Treatments different from those listed above should be carefully considered only when conventional treatments are not effective.²³⁵

Monitoring the possible toxic effect of treatment

Theoretically, the possible toxic effects of treatment should be monitored. However, despite some studies reported possible nephrotoxicity of antimonials,^{85,236} others did not confirm this finding,²³⁷ and recent investigations demonstrated that no toxic effects on heart or pancreas are induced by these drug category in dogs, differently from what is observed in humans.^{238,239} Therefore, toxic

774 effects should be monitored only in selected dogs, particularly when peculiar clinical findings are
775 present or history might lead to hypothesize any drug adversity. The only possible adverse effect of
776 allopurinol is the formation of xanthine crystals, and possibly urolithes, in urine.²⁴⁰ These occur
777 very frequently and may be sometime abundant although associated clinical signs and urolith
778 formation are less common and suspension of treatment is unusual. Therefore, the analysis of urine
779 sediment should be always included in the laboratory workup when allopurinol is included in the
780 treatment protocol or when urine appears macroscopically turbid or forms an evident pellet after
781 centrifugation (figure 8).

782

783 *Monitoring the clinical status*

784 The clinical presentation of leishmaniotic dogs may be extremely variable. Therefore, it is not
785 possible to define, *a priori*, a common and standardized laboratory work up to be used during the
786 follow up. However, two main aspects need to be monitored each time, namely the presence of
787 renal disease and inflammation.

788 Renal function should be evaluated through the analysis of serum concentrations of creatinine and,
789 especially, through sequential quantification of proteinuria, due to its role as a risk factor for the
790 progression of CKD.¹³⁹ Proteinuria has been recently shown to be a negative prognostic factor in
791 leishmaniotic dogs.¹²¹ After conventional leishmanicidal treatment, the degree of proteinuria
792 decreases in 4-8 weeks,²⁴¹ thus, additional pharmacological treatments for proteinuria should be
793 decided thereafter. The possibility to restore normal renal function depends on the severity of renal
794 damage at the time of first diagnosis. Therefore, creatinine and proteinuria should be repeatedly
795 assessed during the follow up. The frequency of testing depends on the severity of CKD: dogs in
796 IRIS stages 3 or 4⁸⁹ should be frequently tested also during the treatment period. Conversely, dogs
797 in IRIS stages 1 or 2⁸⁹ should be tested at the end of the first treatment cycle and then every 6-12
798 months.²⁴² This latter approach is also recommended by the IRIS Glomerular Disease Study
799 Group,²⁴³ that suggests to evaluate renal functions one month after the end of leishmanicidal

800 treatment, and to decide the frequency of further testing depending on the post-treatment IRIS
801 stage: specifically, tests should be run after 12 months in stage I dogs, every 6 months in dogs in
802 stage 2, every 3 months in dogs in stage 3 and every 6 weeks in dogs in stage 4.

803 The inflammatory status may be monitored through sequential analysis of electrophoretograms and
804 of acute phase proteins, whereas the simple evaluation of total protein, albumin or A/G ratio, may
805 not be helpful because it is very likely that, despite treatment decreases globulin concentrations,
806 albumin concentrations remain low in dogs with persistent glomerular damage and proteinuria, in
807 turn leading to only minor changes in the A/G ratio. Differently, serum protein electrophoresis
808 allow to detect a progressive decrease of α - and γ -globulins. These decreases start to become evident
809 after 2-3 weeks and 4-6 weeks, respectively, following treatment with antimonials.²⁴⁴ Therefore, the
810 first useful electrophoretogram to monitor the efficiency of treatment should be run not earlier than
811 one month after treatment begin.²⁴² The complete normalization of electrophoretograms, however,
812 requires at least 90-120 days.²⁴⁰ If after 2-3 months the electrophoretograms still show abnormal
813 profiles, the possible presence of concurrent diseases such as other vector-borne diseases should be
814 considered, especially if the gammopathy tends to be characterized by narrower peaks (see figure
815 4). Treatments with miltefosine or with other drugs may require longer times to be beneficial (more
816 than 2 months to observe a decrease in γ -globulins) and are also characterized by more frequent of
817 relapses after transient normalization of laboratory profiles.^{245,246} Compared with serum protein
818 electrophoresis, monitoring the concentration of APPs provides earlier information regarding the
819 success of treatments with antimonials. CRP and SAA start to decrease in two weeks after treatment
820 and may return within the reference intervals in about one month.^{131,132,244} The normalization of
821 PON-1 and HDL is even more rapid: significant increases may be observed 3-7 days after treatment
822 and values return within the reference intervals in two weeks.^{112,244} Therefore, to assess the efficacy
823 of treatment, it may be advisable to measure the serum activity of PON-1 or the concentration of
824 HDLs or APPs 1-2 weeks after the first administration of drugs, when other clinical or
825 clinicopathological changes are likely still abnormal.

826

827 *Monitoring the parasitological status*

828 As at first diagnosis, the parasitological status can be monitored indirectly, through the assessment
829 of antibody titers, or by direct evaluation of the parasite presence.

830 In case of successful treatment, a decrease of antibody titers may be expected over time; serology
831 can be repeated during the follow up. However, antibody titers tend to decrease at least 6 month
832 after treatment with antimonials combined with continuous allopurinol administration.²⁴⁰ Hence,
833 serology should not be included in the first laboratory tests performed shortly after treatment to
834 monitor the clinical status. Antibody titers should be determined during subsequent evaluations.²⁴²
835 With regard to serological results, it should be kept in mind that a complete negativization of
836 antileishmanial antibodies is unlikely, especially for dogs living in endemic areas that may be
837 repeatedly exposed to the parasite, boosting the antibody response. Therefore, sequential serological
838 tests during the follow up should aim to assess whether antibody titers decrease to values consistent
839 with the simple exposure (i.e. to less than 4 folds the threshold value of the laboratory).¹⁵

840 In order to assess whether treatment completely eradicates the infection, ideally the presence of
841 parasites should be assessed in the tissues in which the parasite may establish a latent infection and
842 using very sensitive techniques. For this purpose the residual parasites burden should be evaluated
843 with repeated quantitative PCR analyses on bone marrow, spleen or lymph nodes, if still palpable.¹⁵
844 However this procedure is invasive and it is difficult that owners accept the analysis, especially if
845 treatment has been successful and the dog is clinically healthy. Therefore, in routine practice the
846 evaluation of treatment efficacy is usually assessed with serology or quantitative PCR analysis in
847 blood. If treatment is successful, the latter test may show a clear decrease of the *Leishmania* DNA
848 copies after 3 to 6 months of therapy, with complete negativization occurring between 6 and 12
849 months.¹⁸²

850

851 *Conclusive remarks and recommended protocols*

852 Diagnosing leishmaniasis in dogs may be difficult due to the complex pathogenesis and broad
853 spectrum of clinical and clinicopathological findings. Hence tests that need to be included in the
854 diagnostic protocol vary in each case or epidemiological scenario.²⁴⁷

855 In clinically healthy dogs living or having travelled to an endemic area serology may be the test of
856 choice to assess any possible exposure to parasites. Based on the median time to achieve
857 seroconversion,¹⁷⁶ serology should be performed at least 6 months after exposure, or periodically
858 (e.g. every 6 months-1 year) in dogs living in endemic areas. If serology is positive, it is important
859 to quantify the antibody response: a low antibody titer may be consistent with exposure or an early
860 phase of infection, while a high antibody titer may suggest infection or disease.^{14,15} Therefore, the
861 subsequent diagnostic steps should confirm the suspected infection through cytological and PCR
862 analysis of sensitive tissues, and/or identification of possible clinical or laboratory alterations,
863 especially in dogs with high antibody titers. If serology or PCR is positive and tests have been
864 performed beyond the transmission season, the laboratory workup should identify the most common
865 abnormalities of dogs with leishmaniasis but no clinical sign (e.g. anemia, abnormal serum protein
866 electrophoresis, proteinuria). If changes are detected, additional clinical or laboratory tests must be
867 performed in order to stage the disease (e.g. tests recommended by the IRIS guidelines for CKD,⁸⁹
868 tests to quantify the acute phase response of inflammation).

869 If the dog is examined because clinical abnormalities are evident, the veterinarian should try to
870 sample any accessible lesion to obtain cytological smears.¹⁵ If amastigotes are documented and the
871 cytological pattern is consistent with leishmaniasis the dog is considered sick; thus, next diagnostic
872 steps should clarify whether a systemic involvement is also present (e.g. hematological disorder,
873 inflammation, nephropathy) and quantify the antibody titer or parasite burden with quantitative
874 PCR to obtain baseline values useful during treatment follow-up. Conversely, if amastigotes are not
875 observed but cytological patterns are consistent with leishmaniasis, the lesion can be further
876 analyzed by histology combined with immunohistochemistry, in situ hybridization or PCR.^{14,15} A
877 positive result with one of these additional tests should lead to investigate further the general health

878 status of the sick dog; conversely, if tests are negative, the presence of infection should be assessed
879 in the bone marrow through cytology and/or PCR and, in case of positive results, further
880 clinopathological tests should be performed as above.^{14,15} Regarding serology, its use is advisable
881 to confirm the disease in dogs suspected to have leishmaniasis but yielding no parasites at
882 cytological examination. Higher titers are more likely to occur in dogs with leishmaniasis and a
883 more severe clinical presentation.

884

885 **Conflict of interest statement**

886 The authors are members of the Canine Leishmaniasis Working Group (CLWG). None of
887 the authors of this paper has a financial or personal relationship with other people or organisations
888 that could inappropriately influence or bias the content of the paper.

889

890 **Acknowledgements:**

891 The authors are grateful to Dr. Alessia Giordano for the collaboration in revising the manuscript.

892

893 **References**

- 894 1. Kuhls K, Alam MZ, Cupolillo E, Ferreira GE, et al. Comparative microsatellite typing of
895 New World *Leishmania infantum* reveals low heterogeneity among populations and its
896 recent Old World origin. *PLoS Negl Trop Dis*. 2011;5:e1155.
- 897 2. Rosypal AC, Troy GC, Zajac AM, Frank G, Lindsay DS. Transplacental transmission of a
898 North American isolate of *Leishmania infantum* in an experimentally infected beagle. *J*
899 *Parasitol*. 200;91:970-972.
- 900 3. De Freitas E, Melo MN, da Costa-Val AP, Michalick MSM. Transmission of *Leishmania*
901 *infantum* via blood transfusion in dogs: potential for infection and importance of clinical
902 factors. *Vet Parasitol*. 2006;137:159-167.

4. Silva FL, Oliveira RG, Silva TM, Xavier MN, Nascimento EF, Santos RL. Venereal transmission of canine visceral leishmaniasis. *Vet Parasitol.* 2009;160:55-59.
5. Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L. Phlebotomine sandflies and the spreading of leishmaniasis and other diseases of public health concern. *Med Vet Entomol.* 2013;27:123-147.
6. Bates PA. Transmission of *Leishmania* metacyclic promastigotes by phlebotomine sand flies. *Int J Parasitol.* 2007;37:1097-1106.
7. Sacks D, Noben-Trauth N. The immunology of susceptibility and resistance to *Leishmania major* in mice. *Nat Rev Immunol.* 2002;2:845-858;
8. Engwerda CR, Ato M, Kaye PM. Macrophages, pathology and parasite persistence in experimental visceral leishmaniasis. *Trends Parasitol.* 2004;20:524-530.
9. Oliva G, Scalone A, Foglia Manzillo V, et al. Incidence and time course of *Leishmania infantum* infections examined by parasitological, serologic, and nested-PCR techniques in a cohort of naive dogs exposed to three consecutive transmission seasons. *J Clin Microbiol.* 2006;44:1318-1322.
10. Pinelli E, Killick-Kendrick R, Wagenaar J, Bernadina W, del Real G, Ruitenbergh J. Cellular and humoral immune responses in dogs experimentally and naturally infected with *Leishmania infantum*. *Infect Immun.* 1994;62:229-235.
11. Santos-Gomes GM, Rosa R, Leandro C, Cortes S, Romão P, Silveira H. Cytokine expression during the outcome of canine experimental infection by *Leishmania infantum*. *Vet Immunol Immunopathol.* 2002;88:21-30.
12. Brachelente C, Muller N, Doherr MG, Sattler U, Welle M. Cutaneous leishmaniasis in naturally infected dogs is associated with a T helper-2-biased immune response. *Vet Pathol.* 200;42:166-175.
13. Chamizo C, Moreno J, Alvar J. Semi-quantitative analysis of cytokine expression in asymptomatic canine leishmaniasis. *Vet Immunol Immunopathol.* 2005;103:67-75.

14. Solano-Gallego L, Koutinas A, Miró G, et al. Directions for the diagnosis, clinical staging, treatment and prevention of canine leishmaniosis. *Vet Parasitol.* 2009;165:1-18.
15. Paltrinieri S, Solano-Gallego L, Fondati A, et al. Guidelines for diagnosis and clinical classification of leishmaniasis in dogs. *J Am Vet Med Assoc.* 2010;236:1184-1191.
16. Alvar J, Canavate C, Molina R, Moreno J, Nieto J. Canine leishmaniasis. *Adv Parasitol.* 2004;57:1-88.
17. Fisa R, Gallego M, Castillejo S, et al. Epidemiology of canine leishmaniosis in Catalonia (Spain) the example of the Priorat focus. *Vet Parasitol.* 1999;83:87-97.
18. Cortes S, Vaz Y, Neves R, Maia C, Cardoso L, Campino L. Risk factors for canine leishmaniasis in an endemic Mediterranean region. *Vet Parasitol.* 2012;189:189-96.
19. Slappendel RJ. Canine leishmaniasis. A review based on 95 cases in The Netherlands. *Vet Q.* 1988;10:1-16.
20. Ciaramella P, Oliva G, Luna RD, et al. A retrospective clinical study of canine leishmaniasis in 150 dogs naturally infected by *Leishmania infantum*. *Vet Rec.* 1997;141:539-543.
21. Koutinas AF, Polizopoulou ZS, Saridomichelakis MN, Argyriadis D, Fytianou A, Plevraki KG. Clinical consideration on canine leishmaniasis in Greece: a retrospective study of 158 cases (1989-1996). *J Am Anim Hosp Assoc.* 1999;35:376-383.
22. Noli C, Saridomichelakis MN. An update on the diagnosis and treatment of canine leishmaniosis caused by *Leishmania infantum* (syn. *L. chagasi*). *Vet J.* 2014;202:425-435.
23. Naranjo C, Fondevila D, Leiva M, Roura X, Peña T. Detection of *Leishmania spp.* and associated inflammation in ocular-associated smooth and striated muscles in dogs with patent leishmaniosis. *Vet Ophthalmol.* 2010;13:139-143.
24. Peña MT, Naranjo C, Klauss G, et al. Histopathological features of ocular leishmaniosis in the dog. *J Comp Pathol.* 2008;138:32-39.
25. Sbrana S, Marchetti V, Mancianti F, Guidi G, Bennett D. Retrospective study of 14 cases of canine arthritis secondary to *Leishmania* infection. *J Small Anim Pract.* 2014;55:309-13.

26. Santos M, Marcos R, Assuncao M, Matos AJ. Polyarthrititis associated with visceral leishmaniasis in a juvenile dog, *Vet Parasitol.* 2006;141:340-344.
27. Foglia Manzillo V, Di Muccio T, Cappiello S, et al. Prospective study on the incidence and progression of clinical signs in naïve dogs naturally infected by *Leishmania infantum*. *PLoS Negl Trop Dis.* 2013;;7:e2225.
28. Adamama-Moraitou KK, Rallis TS, Koytinas AF, Tontis D, Plevraki K, Kritsepi M. Asymptomatic colitis in naturally infected dogs with *Leishmania infantum*: a prospective study. *Am J Trop Med Hyg.* 2007;76:53-57.
29. Parpaglia ML, Vercelli A, Cocco R, Zobba R, Manunta ML. Nodular lesions of the tongue in canine leishmaniosis. *J Vet Med A Physiol Pathol Clin Med.* 2007;54:414-417.
30. Silva FL, Rodrigues AA, Rego IO, et al. Genital lesions and distribution of amastigotes in bitches naturally infected with *Leishmania chagasi*. *Vet Parasitol.* 2008;151:86-90.
31. Silva JF, Guimarães LB, Ribeiro LR, Moreira MV, Serakides R, Ocarino NM. Acute oesophageal necrosis concurrent with *Leishmania chagasi* infection in a dog. *J Comp Pathol.* 2014;;150:148-150.
32. Blavier A, Keroack S, Denerolle P, et al. Atypical forms of canine leishmaniosis. *Vet J.* 2001;162:108-120.
33. Viegas C, Requicha J, Albuquerque C, et al. Tongue nodules in canine leishmaniosis--a case report. *Parasit Vectors.* 2012;5:120.
34. Mir F, Fontaine E, Reyes-Gomez E, Carlus M, Fontbonne A. Subclinical leishmaniasis associated with infertility and chronic prostatitis in a dog. *J Small Anim Pract.* 2012;53:419-422.
35. Ruiz G, Laloy E, Benchekroun G. Chronic gastritis and enterocolitis associated with *Leishmania* infection in an 18-month-old, intact female dog. *Vet Q.* 2015; June 8:1-4 [Epub ahead of print].

36. dos Santos JP, Alves LC, Ramos RA, et al. Histological changes and immunolabeling of *Leishmania infantum* in kidneys and urinary bladder of dogs. *Rev Bras Parasitol Vet.* 2013;22:420-423.
37. Groulade P . Canine leishmaniasis. Clinical haematology and biology of leishmaniasis . *Anim Compagn* 1977;12:121-128.
38. Kiral, FK , Seyrek, K, Pasa, S, Ertabaklar, H, Unsal, C. Some haematological, biochemical and electrophoretic findings in dogs with visceral leishmaniasis. *Revue Med Vet.* 2004;155:226-229.
39. Schultze AE. Interpretation of Canine Leukocyte Response.In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:321-324
40. Nicolato R de C, de Abreu RT, Roatt MB et al. Clinical forms of canine visceral Leishmaniasis in naturally *Leishmania infantum*-infected dogs and related myelogram and hemogram changes. *PLoS One.* 2013;8:e82947.
41. Ruiz de Gopegui, R, Espada Y. Peripheral Blood and Abdominal Fluid from a Dog with Abdominal Distention. *Vet Clin Pathol.* 1998;27:64-67.
42. De Tommasi AS, Otranto D, Furlanello T, et al. Evaluation of blood and bone marrow in selected canine vector-borne diseases. *Parasit Vectors.* 2014;7:534
43. Giudice E, Passantino A Detection of *Leishmania* amastigotes in peripheral blood from four dogs. *Acta Vet Hung.* 2011;59:205-13.
44. Gavazza A , Lubas G , Gugliucci B , Pasquini A, Mancianti F. Hemogram and bone marrow patterns in canine leishmaniasis. *Vet Clin Pathol.* 2002;31:198.
45. Wardrop KJ, Infectious injury to bone marrow In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:118-120.
46. Fry MM. Anemia of nflammatory, neoplastic, renal, and endocrine diseases In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:246-250.

47. Weiss DJ. Myelodysplastic syndromes In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:467-474.
48. Day MJ. Immune - mediated anemias in the dog In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:216-225
49. Tvedten H. Laboratory and clinical diagnosis of anemia In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:152-161.
50. Smith BE, Tompkins MB, Breitschwerdt EB. Antinuclear antibodies can be detected in dog sera reactive to *Bartonella vinsonii subsp. berkhoffii*, *Ehrlichia canis*, or *Leishmania infantum* antigens. *J Vet Intern Med*. 2004;18:47-51.
51. Karagianni AE, Solano-Gallego L, Breitschwerdt EB, et al. Perinuclear antineutrophil cytoplasmic autoantibodies in dogs infected with various vector-borne pathogens and in dogs with immune-mediated hemolytic anemia. *Am J Vet Res*. 2012;73:1403-1409.
52. Terrazzano G, Cortese L, Piantedosi D, et al. Presence of anti - platelet IgM and IgG antibodies in dogs naturally infected by *Leishmania infantum*. *Vet Immunol Immunopathol*. 2006;110:331-337.
53. Cortese L, Sica M, Piantedosi D, et al.: Secondary immune-mediated thrombocytopenia in dogs naturally infected by *Leishmania infantum*. *Vet Rec*. 2009;164:778-782.
54. Cortese L, Terrazzano G, Piantedosi D, et al. Prevalence of anti-platelet antibodies in dogs naturally co-infected by *Leishmania infantum* and *Ehrlichia canis*. *Vet J*. 2011;188:118-21.
55. Green RA, Kabal AL. Hypercoagulable state in three dogs with nephrotic syndrome: role of acquired antithrombin III deficiency. *J Am Vet Med Assoc*. 1982;181:914-917.
56. Honse CO, Figueiredo FB, de Alencar NX, Madeira M de F, Gremião ID, Schubach TM. Disseminated intravascular coagulation in a dog naturally infected by *Leishmania (Leishmania) chagasi* from Rio de Janeiro - Brazil. *BMC Vet Res*. 2013;9:43.
57. Abid M, Kalbantner K, Mischke R. Platelet function in dogs with bacterial infections and leishmaniasis. *Berl Munch Tierarztl Wochenschr*. 2015;128:289-296.

58. Papadogiannakis E, Andritsos G, Kontos V, Spanakos G, Koutis C, Velonakis E.
Determination of CD4+ and CD8+ T cells in the peripheral blood of dogs with
leishmaniosis before and after prolonged allopurinol monotherapy. *Vet J.* 2010;186:262-
263.
59. Rosypal AC, Gogal RM Jr, Zajac AM, Troy GC, Lindsay DS. Flow cytometric analysis of
cellular immune responses in dogs experimentally infected with a North American isolate of
Leishmania infantum. *Vet Parasitol.* 2005;131:45-51.
60. Saridomichelakis MN, Mylonakis ME, Leontides LS, Koutinas AF, Billinis C, Kontos VI.
Evaluation of lymph node and bone marrow cytology in the diagnosis of canine
leishmaniasis (*Leishmania infantum*) in symptomatic and asymptomatic dogs. *Am J Trop
Med Hyg.* 2005;73:82-86.
61. Momo C, Jacintho AP, Moreira PR, Munari DP, Machado GF, Vasconcelos Rde O.
Morphological changes in the bone marrow of the dogs with visceral leishmaniasis. *Vet Med
Int.* 2014;2014:150582.
62. Trópia de Abreu R, Carvalho Md, Carneiro CM, et al. Influence of clinical status and
parasite load on erythropoiesis and leucopoiesis in dogs naturally infected with *leishmania*
(*Leishmania*) *chagasi*. *PLoS One.* 2011;6:e18873.
63. Stockham SL, Scott MA. Bone marrow and lymph node. In: Stockham SL, Scott MA, eds.
Fundamentals of veterinary clinical pathology, 2nd ed. Ames, IA: Blackwell Publishing;
2008:323-368.
64. Weiss DJ . Chronic Inflammation and secondary myelofibrosis In: Weiss DJ, Wardrop KY,
eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:112-117
65. Foglia Manzillo V , Restucci B , Pagano A , Gradoni L, Oliva G. Pathological changes in
the bone marrow of dogs with leishmaniasis. *Vet Rec.* 2006;158:690-694.

66. Lubas G., Caldin M., Wiinberg B., Kristensen A.M. Laboratory testing of coagulation disorders In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:1082-1100.
67. Okano S, Yoshida M, Fukushima U, Higuchi S, Takase K, Hagio M. Usefulness of systemic inflammatory response syndrome criteria as an index for prognosis judgement. *Vet Rec*. 2002;150:245-246.
68. Braun JP, Lefebvre, HP. Kidney function and damage. In: Kaneko JJ, Harvey JW, Bruss ML.eds. *Clinical Biochemistry of Domestic Animals*, 6th ed. San Diego, CA: Elsevier; 2008:457-528.
69. Borgatti A. Plasma cell tumors In: Weiss DJ, Wardrop KY, eds. *Schalm's Veterinary Hematology*, 6th ed. Ames, IA: Wiley-Blackwell; 2010:511-519
70. Lennon EM, Hanel RM, Walker JM, Vaden SL. Hypercoagulability in dogs with protein-losing nephropathy as assessed by thromboelastography. *J Vet Intern Med*. 2013;27:462-468.
71. Bruno B, Maurella C, Falco S, et al. Thromboelastometric evaluation of hemostasis in dogs infected with *Leishmania infantum*. *J Vet Emerg Crit Care (San Antonio)*. 2015;25:502-511.
72. Brooks AC, Guillaumin J, Cooper ES, Couto CG. Effects of hematocrit and red blood cell-independent viscosity on canine thromboelastographic tracings. *Transfusion*. 2014;54:727-734.
73. McMichael MA, Smith SA, Galligan A, Swanson KS. In vitro hypercoagulability on whole blood thromboelastometry associated with in vivo reduction of circulating red cell mass in dogs. *Vet Clin Pathol*. 2014;43:154-163.
74. Vamvakidis CD1, Koutinas AF, Kanakoudis G, Georgiadis G, Saridomichelakis M. Masticatory and skeletal muscle myositis in canine leishmaniasis (*Leishmania infantum*). *Vet Rec*. 2000;146:698-703.

75. Paltrinieri S., Cazzaniga S., Pinto da Cunha N., Giordano A. Electrophoretic fractionation of creatine kinase isoenzymes and macroenzymes in clinically healthy dogs and cats and preliminary evaluation in central neurologic diseases. *Vet Clin Pathol.* 2010;39:329-333.
76. Grano FG, Melo GD, Belinchón-Lorenzo S, Gómez-Nieto LC, Machado GF. First detection of *Leishmania infantum* DNA within the brain of naturally infected dogs. *Vet Parasitol.* 2014;204:376-380.
77. Márquez M1, Pedregosa JR, López J, Marco-Salazar P, Fondevila D, Pumarola M. *Leishmania* amastigotes in the central nervous system of a naturally infected dog. *J Vet Diagn Invest.* 2013;25:142-146.
78. José-López R1, la Fuente CD, Añor S. Presumed brain infarctions in two dogs with systemic leishmaniasis. *J Small Anim Pract.* 2012;53:554-557.
79. Silvestrini P, Piviani M, Alberola J, et al. Serum cardiac troponin I concentrations in dogs with leishmaniasis: correlation with age and clinicopathologic abnormalities. *Vet Clin Pathol.* 2012;41:568–574.
80. Rosa FA1, Leite JH, Braga ET, et al. Cardiac lesions in 30 dogs naturally infected with *Leishmania infantum chagasi*. *Vet Pathol.* 2014;51:603-606.
81. Alves GB, Pinho FA, Silva SM, Cruz MS, Costa FA. Cardiac and pulmonary alterations in symptomatic and asymptomatic dogs infected naturally with *Leishmania (Leishmania) chagasi*. *Braz J Med Biol Res.* 2010;43:310-315.
82. Momo C, Rocha NA, Moreira PR, et al. Morphological changes and parasite load of the adrenal from dogs with visceral leishmaniasis. *Rev Bras Parasitol Vet.* 2014;23:30-35.
83. Saridomichelakis MN, Xenoulis PG, Chatzis MK, et al. Thyroid function in 36 dogs with leishmaniosis due to *Leishmania infantum* before and during treatment with allopurinol with or without meglumine antimonate. *Vet Parasitol.* 2013;197:22-28.
84. Zatelli A, Borgarelli M, Santilli R, et al. Glomerular lesions in dogs infected with *Leishmania* organisms. *Am J Vet Res.* 2003;64:558-561.

85. Aresu L, Benali S, Ferro S, et al. Light and electron microscopic analysis of consecutive renal biopsy specimens from *Leishmania*-seropositive dogs. *Vet Pathol*. 2013;50:753-760.
86. Rigo RS, Carvalho CM, Honer MR, et al. Renal histopathological findings in dogs with visceral leishmaniasis. *Rev Inst Med Trop Sao Paulo*. 2013;55:113-116.
87. Koutinas AF, Koutinas CK. Pathologic mechanisms underlying the clinical findings in canine leishmaniasis due to *Leishmania infantum/chagasi*. *Vet Pathol*. 2014;51:527-538.
88. Braga ET, Leite JH, Rosa FA, et al. Hypertension and its correlation with renal lesions in dogs with leishmaniosis. *Rev Bras Parasitol Vet*. 2015;24:45-51.
89. International renal Interest Society (IRIS): Guidelines for staging chronic kidney disease (CKD). Available at: www.iris-kidney.com. Accessed, October 20, 2015
90. Brown S, Atkins C, Bagley R, et al. Guidelines for the identification, evaluation, and management of systemic hypertension in dogs and cats. *J Vet Intern Med*. 2007;21:542-558.
91. Ribeiro RR, da Silva SM, Fulgêncio Gde O, Michalick MS, Frézard FJ. Relationship between clinical and pathological signs and severity of canine leishmaniasis. *Rev Bras Parasitol Vet*. 2013;22:373-378.
92. Braun JP, Lefebvre HP, Watson AD. Creatinine in the dog: a review. *Vet Clin Pathol*. 2003;32:162-179.
93. Stockham SL, Scott MA. Urinary system. In: Stockham SL, Scott MA, eds. *Fundamentals of veterinary clinical pathology*, 2nd ed. Ames, IA: Blackwell Publishing; 2008:415-494.
94. Von Hendy-Willson VE, Pressler BM. An overview of glomerular filtration rate testing in dogs and cats. *Vet J*. 2011;188:156-165.
95. Sasaki Y, Iwama R, Shimamura S, et al. Estimation of glomerular filtration rate in dogs by a single-blood sample method involving iodixanol. *Am J Vet Res*. 2015;76:828-835.
96. Jensen AL, Bomholt M, Moe L. Preliminary evaluation of a particle-enhanced turbidimetric immunoassay (PETIA) for the determination of serum cystatin C-like immunoreactivity in dogs. *Vet Clin Pathol*. 2001;30:86-90

97. Almy FS, Christopher MM, King DP, Brown SA . Evaluation of cystatin C as an endogenous marker of glomerular filtration rate in dogs. *J Vet Intern Med.* 2002;16:45-51.
98. Miyagawa Y, Takemura N, Hirose H. Evaluation of the measurement of serum cystatin C by an enzyme-linked immunosorbent assay for humans as a marker of the glomerular filtration rate in dogs. *J Vet Med Sci.* 2009;71:1169-1176.
99. Ghys L, Paepe D, Smets P, Lefebvre H, Delanghe J, Daminet S. Cystatin C: a new renal marker and its potential use in small animal medicine. *J Vet Intern Med.* 2014;28:1152-1164.
100. Pasa S, Bayramli G, Atasoy A, Karul A, Ertug S, Ozensoy Toz S. Evaluation of serum cystatin-C in dogs with visceral leishmaniasis. *Vet Res Commun.* 2009;33:529-534.
101. Monti P, Benckekroun G, Berlato D, Archer J. Initial evaluation of canine urinary cystatin C as a marker of renal tubular function. *J Small Anim Pract.* 2012;53:254-259.
102. García-Martínez JD, Martínez-Subiela S, Tvarijonaviciute A, Caldin M, Ceron JJ. Urinary ferritin and cystatin C concentrations at different stages of kidney disease in leishmaniotic dogs. *Res Vet Sci.* 2015;99:204-207.
103. Jepson RE, Syme HM, Vallance C, Elliott J., Plasma asymmetric dimethylarginine, symmetric dimethylarginine, l-arginine, and nitrite/nitrate concentrations in cats with chronic kidney disease and hypertension. *J Vet Intern Med.* 2008;22:317-324.
104. Nabity MB, Lees GE, Boggess MM, et al. Symmetric Dimethylarginine Assay Validation, Stability, and Evaluation as a Marker for the Early Detection of Chronic Kidney Disease in Dogs. *J Vet Intern Med.* 2015;29:1036-1044.
105. Braff J, Obare E, Yerramilli M, Elliott J, Yerramilli M. Relationship between serum symmetric dimethylarginine concentration and glomerular filtration rate in cats. *J Vet Intern Med.* 2014;28:1699-1701.
106. Hall JA, Yerramilli M, Obare E, Yerramilli M, Yu S, Jewell DE. Comparison of serum concentrations of symmetric dimethylarginine and creatinine as kidney function

1159 biomarkers in healthy geriatric cats fed reduced protein foods enriched with fish oil, L-
 1160 carnitine, and medium-chain triglycerides. *Vet J.* 2014;202:588-596.

1161 107. Yerramilli M, Yerramilli M, Obare E, Jewell DE, Hall JA. Symmetric
 1162 dimethylarginine (SDMA) increases earlier than serum creatinine in dogs with chronic
 1163 kidney disease (CKD). *J Vet Intern Med.* 2014;28:1084–1085.

1164 108. Hall JA, Yerramilli M, Obare E, Yerramilli M, Melendez LD, Jewell DE.
 1165 Relationship between lean body mass and serum renal biomarkers in healthy dogs. *J Vet*
 1166 *Intern Med.* 2015;29:808-814.

1167 109. Stanger, O., Herrmann, W., Pietrzik, K., et al. DACH-LIGA Homocystein (German,
 1168 Austrian and Swiss Homocysteine Society): Consensus paper on the rational clinical use of
 1169 homocysteine, folic acid and B-vitamins in cardiovascular and thrombotic diseases:
 1170 Guidelines and recommendations. *Clin Chem Lab Med.* 2003;41:1392-1403.

1171 110. Schiffrin EL. Role of endothelin-1 in hypertension. *J Am Heart Assoc.* 1999;34:876-
 1172 881.

1173 111. Barton M, Yanagisawa M. Endothelin: 20 years from discovery to therapy. *Can J*
 1174 *Physiol Pharmacol.* 2008;86:485-498.

1175 112. Ceron JJ, Eckersall PD, Martinez-Subiela S. Acute phase proteins in dogs and cats:
 1176 Current knowledge and future perspectives. *Vet Clin Pathol.* 2005;34:85-99.

1177 113. Raila J, Schweigert FJ, Kohn B. C-reactive protein concentrations in serum of dogs
 1178 with naturally occurring renal disease. *J Vet Diagn Invest.* 2011;23:710-715.

1179 114. Rossi S, Rossi G, Giordano A, Paltrinieri S. Homocysteine measurement by an
 1180 enzymatic method and potential role of homocysteine as a biomarker in dogs. *J Vet Diagn*
 1181 *Invest.* 2008;20:644-649.

1182 115. Rossi G, Giordano A, Breda S, et al. Big-endothelin 1 (Big ET-1) and homocysteine
 1183 in the serum of dogs with chronic kidney disease. *Vet J.* 2013;198:109-115.

116. Martínez-Subiela S, García-Martínez JD, Tvarijonaviciute A, et al. Urinary C reactive protein levels in dogs with leishmaniasis at different stages of renal damage. *Res Vet Sci.* 2013;95:924-929.
117. Tvarijonaviciute A, Ceron JJ, Martinez-Subiela S, García-Martínez JD. Serum and urinary adiponectin in dogs with renal disease from leishmaniasis. *Vet Rec.* 2012;171:297.
118. Almeida MA, Jesus EE, Sousa-Atta ML, Alves LC, Berne ME, Atta AM. Clinical and serological aspects of visceral leishmaniasis in northeast Brazilian dogs naturally infected with *Leishmania chagasi*. *Vet Parasitol.* 2005;127:227-232.
119. Eckersall PD. Proteins, proteomics, and the dysproteinemias. In: Kaneko JJ, Harvey JW, Bruss ML. eds. *Clinical Biochemistry of Domestic Animals*, 6th ed. San Diego, CA: Elsevier; 2008:117-156
120. Proverbio D, Spada E, Bagnagatti de Giorgi G, Perego R, Valena E. Relationship between *Leishmania* IFAT titer and clinicopathological manifestations (clinical score) in dogs. *Biomed Res Int.* 2014;2014:412808.
121. Geisweid K, Mueller R, Sauter-Louis C, Hartmann K. Prognostic analytes in dogs with *Leishmania infantum* infection living in a non-endemic area. *Vet Rec.* 2012;171:399.
122. Vailati Facchini R, Bertazzolo W, Zuliani D, et al. Detection of biclonal gammopathy by capillary zone electrophoresis in a cat and a dog with plasma cell neoplasia. *Vet Clin Pathol.* 2010;39:440-446.
123. Font A , Closa JM , Mascort J . Monoclonal gammopathy in a dog with visceral leishmaniasis. *J Vet Int Med.* 1994;8:233-235.
124. Giordano A, Paltrinieri S. Interpretation of capillary zone electrophoresis compared with cellulose acetate and agarose gel electrophoresis: reference intervals and diagnostic efficiency in dogs and cats. *Vet Clin Pathol.* 2010;39:464-473.

- 1208 125. Antognoni MT, Biretoni F, Miglio A, Lalli P, Porciello F, Mangili Pecci V.
1209 Monoclonal gammopathy associated with multiple myeloma and visceral leishmaniasis in
1210 the dog: a comparison of two cases. *Vet Res Commun*. 2010;34 Suppl 1:S97-101.
- 1211 126. Geigy C, Riond B, Bley CR, Grest P, Kircher P, Lutz H. Multiple myeloma in a dog
1212 with multiple concurrent infectious diseases and persistent polyclonal gammopathy. *Vet Clin*
1213 *Pathol*. 2013;42:47-54.
- 1214 127. Martínez-Subiela S, Tecles F, Eckersall PD, Cerón JJ. Serum concentrations of acute
1215 phase proteins in dogs with leishmaniasis. *Vet Rec*. 2002;150:241-244.
- 1216 128. Martinez-Subiela S, Strauss-Ayali D, Cerón JJ, Baneth G. Acute phase protein
1217 response in experimental canine leishmaniasis. *Vet Parasitol*. 2011;180:197-202.
- 1218 129. Silvestrini P, Zoia A, Planellas M, et al. Iron status and C-reactive protein in canine
1219 leishmaniasis. *J Small Anim Pract*. 2014;55:95-101.
- 1220 130. Martinez-Subiela S, Cerón JJ, Strauss-Ayali D, et al. Serum ferritin and paraoxonase-
1221 1 in canine leishmaniosis. *Comp Immunol Microbiol Infect Dis*. 2014;37:23-29.
- 1222 131. Martínez-Subiela S, Bernal LJ, Cerón JJ. Serum concentrations of acute-phase
1223 proteins in dogs with leishmaniosis during short-term treatment. *Am J Vet Res*.
1224 2003;64:1021-1026.
- 1225 132. Sasanelli M, Paradies P, de Caprariis D, et al. Acute-phase proteins in dogs naturally
1226 infected with *Leishmania infantum* during and after long-term therapy with allopurinol. *Vet*
1227 *Res Commun*. 2007;31 Suppl 1:335-338.
- 1228 133. Rossi G, Giordano A, Pezzia F, Kjelgaard-Hansen M, Paltrinieri S. Serum
1229 paraoxonase activity (PON1) in dogs: preanalytical and analytical factors and correlation
1230 with C-reactive protein and alpha-2-globulin. *Vet Clin Pathol*. 2013;42:329-341.
- 1231 134. Ibba F, Rossi G, Meazzi S, Giordano A, Paltrinieri S. Serum concentration of high
1232 density lipoproteins (HDLs) in leishmaniotic dogs. *Res Vet Sci*. 2015;98:89-91.

- 1233 135. Tonin AA, Calado AM, Bottari NB, et al. Novel markers of inflammatory response
1234 and hepatic dysfunction in canine leishmaniasis. *Comp Immunol Microbiol Infect Dis*. 2015
1235 pii: S0147-9571(15)00064-8. doi: 10.1016/j.cimid.2015.09.004. [Epub ahead of print]
- 1236 136. Osborne CA, Stevens J. Urine sediment. In: Osborne CA, Stevens J. *Urinalysis: a*
1237 *clinical guide to compassionate patient care*. Shawnee, KS; Bayer Corporation: 1999;125-
1238 150.
- 1239 137. Heuter KJ, Buffington CA, Chew DJ. Agreement between two methods for
1240 measuring urine pH in cats and dogs. *J Am Vet Med Assoc*. 1998;213:996-998.
- 1241 138. Vaden SL, Pressler BM, Lappin MR, Jensen WA. Effects of urinary tract
1242 inflammation and sample blood contamination on urine albumin and total protein
1243 concentrations in canine urine samples. *Vet Clin Pathol*. 2004;33:14-19.
- 1244 139. Jacob F, Polzin DJ, Osborne CA, et al. Evaluation of the association between initial
1245 proteinuria and morbidity rate or death in dogs with naturally occurring chronic renal
1246 failure. *J Am Vet Med Assoc*. 2005;226:393-400.
- 1247 140. Lees GE, Brown SA, Elliott J, Grauer GE, Vaden SL; American College of
1248 Veterinary Internal Medicine. Assessment and management of proteinuria in dogs and cats:
1249 2004 ACVIM Forum Consensus Statement (small animal). *J Vet Intern Med*. 2005;19:377-
1250 385.
- 1251 141. Beatrice L, Nizi F, Callegari D, et al. Comparison of urine protein-to-creatinine ratio
1252 in urine samples collected by cystocentesis versus free catch in dogs. *J Am Vet Med Assoc*.
1253 2010;236:1221-1224.
- 1254 142. Zatelli A, Paltrineiri S, Nizi F, Roura X, Zini E. Evaluation of a urine dipstick test for
1255 confirmation or exclusion of proteinuria in dogs. *Am J Vet Res*. 2010;71:235-240.
- 1256 143. IRIS Canine GN Study Group Diagnosis Subgroup, Littman MP, Daminet S, Grauer
1257 GF, Lees GE, van Dongen AM. Consensus recommendations for the diagnostic

- investigation of dogs with suspected glomerular disease. *J Vet Intern Med.* 2013;;27 Suppl 1:S19-526.
144. Rossi G, Giori L, Campagnola S, Zatelli A, Zini E, Paltrinieri S. Evaluation of factors that affect analytic variability of urine protein-to-creatinine ratio determination in dogs. *Am J Vet Res.* 2012;73:779-788.
 145. Rossi G, Bertazzolo W, Dondi F, et al. Inter-laboratory variability in protein:creatinine (UPC) ratio on canine urine. *Vet J.* 2015;204:66-72.
 146. Rossi G, Bertazzolo W, Binnella M, Scarpa P, Paltrinieri S. Measurement of proteinuria in dogs: analytical and diagnostic differences using two laboratory methods *Vet Clin Pathol.* In press
 147. LeVine DN, Zhang D, Harris T, Vaden SL. The use of pooled vs serial urine samples to measure urine protein:creatinine ratios. *Vet Clin Pathol.* 2010;39:53-56.
 148. Smets PM, Meyer E, Maddens BE, Duchateau L, Daminet S. Urinary markers in healthy young and aged dogs and dogs with chronic kidney disease. *J Vet Intern Med.* 2010;24:65-72.
 149. Schultze,AE, Jensen RK. Sodium dodecyl sulfate polyacrylamide gel electrophoresis of canine urinary proteins for the analysis and differentiation of tubular and glomerular diseases. *Vet Clin Pathol.* 1998;18:93-97.
 150. Brown JS, Nabity MB, Brock R, Cianciolo R, Lees GE. Comparison of urine sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) with renal histological findings and clinicopathologic data in dogs with renal diseases. *Vet Clin Pathol.* 2010;39:556.
 151. Zini E, Bonfanti U, Zatelli A. Diagnostic relevance of qualitative proteinuria evaluated by use of sodium dodecyl sulfate-agarose renal histologic findings in dogs. *Am J Vet Res.* 2004;65:964–971.

- 1283 152. Giori L, Tricomi FM, Zatelli A, Roura X, Paltrinieri S. High-resolution gel
1284 electrophoresis and sodium dodecyl sulphate–agarose gel electrophoresis on urine samples
1285 for qualitative analysis of proteinuria in dogs. *J Vet Diagn Invest.* 2011;23:682-690.
- 1286 153. Nadaud C, Piane L, Theron ML, Palanche F, Trumel C, Lavoue R. 2014. Urine
1287 electrophoretic protein profile of healthy male dogs. *Vet Clin Pathol.* 2005;44:E8–E9
- 1288 154. Theron ML, Trumel C, Piane L, Lucarelli L, Henrion R, Lavoue R. Effects of storage
1289 conditions on canine urine sodium dodecyl sulfate-agarose gel electrophoresis (SDS-AGE).
1290 *Vet Clin Pathol.* 2013;42:34.
- 1291 155. Ibba F, Mangiagalli G, Paltrinieri S. Urinary gamma-glutamyl transferase (GGT) as a
1292 marker of tubular damage in dogs with canine leishmaniasis, using sodium dodecylsulphate
1293 (SDS) electrophoresis as a reference method. *Vet J. in press*
- 1294 156. Bonfanti U, Zini E, Minetti E, Zatelli A. Free light-chain proteinuria and normal
1295 renal histopathology and function in 11 dogs exposed to *Leishmania infantum*, *Ehrlichia*
1296 *canis*, and *Babesia canis*. *J Vet Intern Med.* 2004;18:618-624.
- 1297 157. Brunker JD, Ponzio NM, Payton ME. Indices of urine N-acetyl-beta-D-
1298 glucosaminidase and gamma-glutamyl transpeptidase activities in clinically normal adult
1299 dogs. *Am J Vet Res.* 2009;70:297-301.
- 1300 158. Matteucci E, Giampietro O. To store urinary enzymes: how and how long? *Kidney*
1301 *Int. Suppl.* 1994;47:S58-559.
- 1302 159. Palacio J, Liste F, Gascón M. Enzymuria as an index of renal damage in canine
1303 leishmaniasis. *Vet Rec.* 1997;140:477-480.
- 1304 160. García-Martínez JD, Tvarijonaviciute A, Cerón JJ, Caldin M, Martínez-Subiela S.
1305 Urinary clusterin as a renal marker in dogs. *J Vet Diagn Invest.* 2012;24:301-306.
- 1306 161. Nabity MB, Lees GE, Cianciolo R, Boggess MM, Steiner JM, Suchodolski JS.
1307 Urinary biomarkers of renal disease in dogs with X-linked hereditary nephropathy. *J Vet*
1308 *Intern Med.* 2012;26:282-293.

162. Fuchs TC, Hewitt P. Biomarkers for drug-induced renal damage and nephrotoxicity-an overview for applied toxicology. *AAPS J.* 2011;13:615-631.
163. Sousa S, Cardoso L, Reed SG, et al. Development of a fluorescent based immunosensor for the serodiagnosis of canine leishmaniasis combining immunomagnetic separation and flow cytometry. *PLoS Negl Trop Dis.* 2013;7:e2371.
164. Ker HG, Coura-Vital W, Aguiar-Soares RD, et al. Evaluation of a prototype flow cytometry test for serodiagnosis of canine visceral leishmaniasis. *Clin Vaccine Immunol.* 2013;20:1792-1798.
165. Ramos-Jesus J, Carvalho KA, Fonseca RA, et al. A piezoelectric immunosensor for *Leishmania chagasi* antibodies in canine serum. *Anal Bioanal Chem.* 2011;401:917-925.
166. Akhoundi B, Mohebbi M, Shojaei S, et al. Rapid detection of human and canine visceral leishmaniasis: assessment of a latex agglutination test based on the A2 antigen from amastigote forms of *Leishmania infantum*. *Exp Parasitol.* 2013;133:307-313.
167. Maia C, Campino L. Methods for diagnosis of canine leishmaniasis and immune response to infection. *Vet Parasitol.* 2008;158:274-287.
168. Rosypal AC, Pick LD, Hernandez JO, Lindsay DS. Evaluation of a novel dried blood spot collection device (HemaSpot™) to test blood samples collected from dogs for antibodies to *Leishmania infantum*. *Vet Parasitol.* 2014;205:338-342.
169. EFSA AHAW Panel (EFSA Panel on Animal Health and Welfare), 2015. Scientific Opinion on canine leishmaniosis. *EFSA Journal* 2015;13:4075, 77 pp.
doi:10.2903/j.efsa.2015.4075
170. Grimaldi G Jr, Teva A, Ferreira AL, et al. Evaluation of a novel chromatographic immunoassay based on Dual-Path Platform technology (DPP® CVL rapid test) for the serodiagnosis of canine visceral leishmaniasis. *Trans R Soc Trop Med Hyg.* 2012;106:54-59.
171. Sagols E, Martin V, Claret E, McGahie D, Ciusinier AM, Gueguen S. Evaluation of the humoral immune response after vaccination with LiESP/QA21 (CaniLeish): interest of

Leishmania specific anti-kinesin antibodies detection. *Proceedings of the 2012 BSAVA Congress*, Birmingham, United Kingdom, April 12th-15th, 2012

172. Ferroglio E, Zanet S, Mignone W, Poggi M, Trisciuglio A, Bianciardi P. Evaluation of a rapid device for serological diagnosis of *Leishmania infantum* infection in dogs as an alternative to immunofluorescence assay and Western blotting. *Clin Vaccine Immunol*. 2013;20:657-659.

173. Solano-Gallego L, Villanueva-Saz S, Carbonell M, Trotta M, Furlanello T, Natale A. Serological diagnosis of canine leishmaniosis: comparison of three commercial ELISA tests (Leiscan, ID Screen and *Leishmania* 96), a rapid test (Speed Leish K) and an in-house IFAT. *Parasit Vectors*. 2014;7:111.

174. Gradoni L, Gramiccia M. Leishmaniosis. In *Manual of diagnostic tests and vaccines for terrestrial animals*. World Organisation for Animal Health (OIE); 2008:240-250. (2014 update available at: http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.01.08_LEISHMANIOSIS.pdf; accessed, October 20, 2015)

175. Rodríguez-Cortés A, Ojeda A, Francino O, López-Fuertes L, Timón M, Alberola J. *Leishmania* infection: laboratory diagnosing in the absence of a "gold standard". *Am J Trop Med Hyg*. 2010;82:251-256.

176. Moreno J, Alvar J. Canine leishmaniasis: epidemiological risk and the experimental model. *Trends Parasitol*. 2002;18:399-405.

177. dos-Santos WL, Jesus EE, Paranhos-Silva M, et al. Associations among immunological, parasitological and clinical parameters in canine visceral leishmaniasis: emaciation, spleen parasitism, specific antibodies and leishmanin skin test reaction. *Vet Immunol Immunopathol*. 2008;5:3-4.

178. Andreadou M, Liandris E, Gazouli M, et al. A novel non-amplification assay for the detection of *Leishmania spp.* in clinical samples using gold nanoparticles. *J Microbiol Methods*. 2014;96:56-61.
179. Chaouch M, Mhadhbi M, Adams ER, et al. Development and evaluation of a loop-mediated isothermal amplification assay for rapid detection of *Leishmania infantum* in canine leishmaniasis based on cysteine protease B genes. *Vet Parasitol*. 2013;198:78-84.
180. Gao CH, Ding D, Wang JY, et al. Development of a LAMP assay for detection of *Leishmania infantum* infection in dogs using conjunctival swab samples. *Parasit Vectors*. 2015;8:370.
181. Cortes S, Rolao N, Ramada J, Campino L. PCR as a rapid and sensitive tool in the diagnosis of human and canine leishmaniasis using *Leishmania donovani* s.l.-specific kinetoplastid primers. *Trans R Soc Trop Med Hyg*. 2004;98:12-17.
182. Francino O, Altet L, Sanchez-Robert E, et al.: Advantages of real-time PCR assay for diagnosis and monitoring of canine leishmaniosis. *Vet Parasitol*. 2006;137:214-221.
183. Martínez V, Quilez J, Sanchez A, Roura X, Francino O, Altet L. Canine leishmaniasis: the key points for qPCR result interpretation. *Parasit Vectors*. 2011;4:57.
184. Solcà Mda S, Guedes CE, Nascimento EG, et al. Qualitative and quantitative polymerase chain reaction (PCR) for detection of *Leishmania* in spleen samples from naturally infected dogs. *Vet Parasitol*. 2012;184:133-140.
185. Santos TR, Carreira VS, Ferrari HF, Moreira MA, Luvizotto MC. Comparison of PCR with stained slides of bone marrow and lymph nodes aspirates with suspect diagnosis for leishmaniasis. *Acta Trop*. 2014;140:137-140.
186. Roura X, Fondevila D, Sanchez A, Ferrer L. Detection of *Leishmania* infection in paraffin-embedded skin biopsies of dogs using polymerase chain reaction. *J Vet Diagn Invest*. 1999;11:385-387.

187. Muller N, Zimmermann V, Forster U, Bienz M, Gottstein B, Welle M. PCR-based detection of canine *Leishmania* infections in formalin-fixed and paraffinembedded skin biopsies: elaboration of a protocol for quality assessment of the diagnostic amplification reaction. *Vet Parasitol.* 2003;114:223-229.
188. Gradoni, L. The diagnosis of canine leishmaniasis. In: Killick-Kendrick R, ed. *Canine leishmaniasis: moving towards a solution. Proceedings of the 2nd International leishmaniasis Forum, Sevilla, Spain.* Boxmeer, NL. Intervet International. 2002:7-14.
189. Ramos RA, Ramos CA, Santos EM, et al. Quantification of *Leishmania infantum* DNA in the bone marrow, lymph node and spleen of dogs. *Rev Bras Parasitol Vet.* 2013;22:346-350.
190. Carvalho Ferreira AL, Carregal VM, de Almeida Ferreira S, Leite RS, de Andrade AS. Detection of *Leishmania infantum* in 4 different dog samples by real-time PCR and ITS-1 nested PCR. *Diagn Microbiol Infect Dis.* 2014;78:418-421.
191. Almeida AB, Sousa VR, Gasparetto ND, et al. Canine visceral leishmaniasis: diagnostic approaches based on polymerase chain reaction employing different biological samples. *Diagn Microbiol Infect Dis.* 2013;76:321-324.
192. Reis LE, Coura-Vital W, Roatt BM, et al. Molecular diagnosis of canine visceral leishmaniasis: a comparative study of three methods using skin and spleen from dogs with natural *Leishmania infantum* infection. *Vet Parasitol.* 2013;197:498-503.
193. Ferreira Sde A, Ituassu LT, de Melo MN, de Andrade AS: Evaluation of the conjunctival swab for canine visceral leishmaniasis diagnosis by PCR-hybridization in Minas Gerais State, Brazil. *Vet Parasitol.* 2008;152:257-263.
194. Gramiccia M, Di Muccio T, Fiorentino E, et al.: Longitudinal study on the detection of canine *Leishmania* infections by conjunctival swab analysis and correlation with entomological parameters. *Vet Parasitol.* 2010;171:223-228.

195. Di Muccio T, Veronesi F, Antognoni MT, Onofri A, Piergili Fioretti D, Gramiccia M. Diagnostic value of conjunctival swab sampling associated with nested PCR for different categories of dogs naturally exposed to *Leishmania infantum* infection. *J Clin Microbiol.* 2012;50:2651-2659.
196. Ferreira Sde A, Almeida GG, Silva Sde O, et al. Nasal, oral and ear swabs for canine visceral leishmaniasis diagnosis: new practical approaches for detection of *Leishmania infantum* DNA. *PLoS Negl Trop Dis.* 2013;7:e2150.
197. Madeira MF, Figueiredo FB, Pinto AG, et al.: Parasitological diagnosis of canine visceral leishmaniasis: is intact skin a good target? *Res Vet Sci.* 2009;87:260-262.
198. Diniz SA, Melo MS, Borges AM, et al. Genital lesions associated with visceral leishmaniasis and shedding of *Leishmania spp.* in the semen of naturally infected dogs. *Vet Pathol.* 2005;42:650-658
199. Manna L, Paciello O, Morte RD, Gravino AE. Detection of *Leishmania* parasites in the testis of a dog affected by orchitis: case report. *Parasit Vectors.* 2012;5:216.
200. Levy E, Mylonakis ME, Saridomichelakis MN, Polizopoulou ZS, Psychogios V, Koutinas AF. Nasal and oral masses in a dog. *Vet Clin Pathol.* 2006;35:115-118
201. Caniatti M, da Cunha NP, Avallone G, et al. Diagnostic accuracy of brush cytology in canine chronic intranasal disease. *Vet Clin Pathol.* 2012;41:133-140.
202. Costa MM, Lima WG, Figueiredo MM, Michalick MS, Tafuri WL, Tafuri WL. Cervical, mandibular, and parotid lymph nodes of dogs naturally infected with *Leishmania infantum*: a histopathologic and immunohistochemistry study and its correlation with facial skin lesions. *Vet Pathol.* 2008;45:613-616.
203. Mylonakis ME, Papaioannou N, Saridomichelakis MN, Koutinas AF, Billinis C, Kontos VI. Cytologic patterns of lymphadenopathy in dogs infected with *Leishmania infantum*. *Vet Clin Pathol.* 2005;34:243-247.

204. Fernandes PJ. Synovial fluid analysis. In: Cowell RL, Tyler RD, Meinkoth JH, DeNicola DB,eds. *Diagnostic cytology and hematology of the dog and cat*. 3rd ed. St. Louis, MO: Mosby Elsevier; 2008:193-209.
205. Desnoyers M, Bedard C, Meinkoth JH, Crystal MA. Cerebrospinal fluid analysis In: Cowell RL, Tyler RD, Meinkoth JH, DeNicola DB,eds. *Diagnostic cytology and hematology of the dog and cat*. 3rd ed. St. Louis, MO: Mosby Elsevier; 2008:215-234.
206. Ferro S, Palmieri C, Cavicchioli L, De Zan G, Aresu L, Benali SL. *Leishmania* amastigotes in neoplastic cells of 3 nonhistiocytic canine tumors. *Vet Pathol*. 2013;50:749-752.
207. Kegler K, Habierski A, Hahn K, Amarilla SP, Seehusen F, Baumgärtner W. Vaginal canine transmissible venereal tumour associated with intra-tumoural *Leishmania spp.* amastigotes in an asymptomatic female dog. *J Comp Pathol*. 2013;149:156-161.
208. Marino G, Gaglio G, Zanghi A. Clinicopathological study of canine transmissible venereal tumour in leishmaniotic dogs. *J Small Anim Pract*. 2012;53:323-327.
209. Foglia Manzillo V, Pagano A, Guglielmino R, Gradoni L, Restucci B, Oliva G. Extranodal gammadelta-T-cell lymphoma in a dog with leishmaniasis. *Vet Clin Pathol*. 2008;37:298-301.
210. Catone G, Marino G, Poglayen G, Gramiccia M, Ludovisi A, Zanghi A. Canine transmissible venereal tumour parasitized by *Leishmania infantum*. *Vet Res Commun*. 2003;27:549-553.
211. Albanese F, Poli A, Millanta F, Abramo F. Primary cutaneous extragenital canine transmissible venereal tumour with *Leishmania*-laden neoplastic cells: a further suggestion of histiocytic origin? *Vet Dermatol*. 2002;13:243-246.
212. Toplu N, Aydogan A. An immunohistochemical study in cases with usual and unusual clinicopathological findings of canine visceral leishmaniosis. *Parasitol Res*. 2011;109:1051-1057.

213. Menezes RC, Figueiredo FB, Wise AG, et al. Sensitivity and specificity of in situ hybridization for diagnosis of cutaneous infection by *Leishmania infantum* in dogs. *J Clin Microbiol.* 2013;51:206-211.
214. Dinhopl N, Mostegl MM, Richter B, et al. In situ hybridisation for the detection of *Leishmania* species in paraffin wax-embedded canine tissues using a digoxigenin-labelled oligonucleotide probe. *Vet Rec.* 2011;169:525.
215. Nieto CG, Navarrete I, Habela MA, Serrano F, Redondo E. Pathological changes in kidneys of dogs with natural *Leishmania* infection. *Vet Parasitol.* 1992;45:33-47.
216. Koutinas AF, Carlotti DN, Koutinas C, Papadogiannakis EI, Spanakos GK, Saridomichelakis MN. Claw histopathology and parasitic load in natural cases of canine leishmaniosis associated with *Leishmania infantum*. *Vet Dermatol.* 2010;21:572-577
217. Naranjo C, Fondevila D, Altet L, et al. Evaluation of the presence of *Leishmania spp.* by real-time PCR in the lacrimal glands of dogs with leishmaniosis. *Vet J.* 2012;193:168-173.
218. Figueiredo MM, Moura EP, Costa MM, et al. Histopathological and parasitological investigations of ear healthy skin of dogs naturally and experimentally infected with *Leishmania (Leishmania) chagasi*. *Histol Histopathol.* 2010;25:877-887.
219. Moreira PR, Vieira LM, de Andrade MM, et al. Immune response pattern of the popliteal lymph nodes of dogs with visceral leishmaniasis. *Parasitol Res.* 2010;107:605-613.
220. Pinto AJ, Figueiredo MM, Silva FL, et al. Histopathological and parasitological study of the gastrointestinal tract of dogs naturally infected with *Leishmania infantum*. *Acta Vet Scand.* 2011;53:67.
221. Bardagi M, Fondevila D, Zanna G, Ferrer L. Histopathological differences between canine idiopathic sebaceous adenitis and canine leishmaniosis with sebaceous adenitis. *Vet Dermatol.* 2010;21:159-165.

222. Bongiorno G, Paparcone R, Foglia Manzillo V, Oliva G, Cuisinier AM, Gradoni L. Vaccination with LiESP/QA-21 (CaniLeish®) reduces the intensity of infection in *Phlebotomus perniciosus* fed on *Leishmania infantum* infected dogs--a preliminary xenodiagnosis study. *Vet Parasitol.* 2013;197:691-695.
223. Longoni SS, Sánchez-Moreno M, López JE, Marín C. *Leishmania infantum* secreted iron superoxide dismutase purification and its application to the diagnosis of canine Leishmaniasis. *Comp Immunol Microbiol Infect Dis.* 2013;36:499-506.
224. Britti D, Gaspari M, Massimini G, Casalnuovo F, Morittu VM, Cuda G. Proteomic analysis in canine leishmaniasis. *Vet Res Commun.* 2010;34 Suppl 1:S91-596.
225. Menezes-Souza D, Corrêa-Oliveira R, Guerra-Sá R, et al. Cytokine and transcription factor profiles in the skin of dogs naturally infected by *Leishmania (Leishmania) chagasi* presenting distinct cutaneous parasite density and clinical status. *Vet Parasitol.* 2011;177:39-49.
226. Di Loria A, Squillacioti C, De Luca A, et al. Increased leptin mRNA expression in the blood of dogs naturally infected by *Leishmania infantum*. *Vet J.* 2014;202:634-636.
227. dos Santos FR, Vieira PM, Correa-Oliveira R, et al. Qualitative and quantitative immunohistochemical evaluation of iNOS expression in the spleen of dogs naturally infected with *Leishmania chagasi*. *Parasitol Res.* 2011;108:1397-1403.
228. Melo GD, Marangoni NR, Marcondes M, Lima VM, Machado GF. High levels of serum matrix metalloproteinases in dogs with natural visceral leishmaniosis: a preliminary report. *Vet J.* 2011;188:243-245.
229. Marangoni NR, Melo GD, Moraes OC, Souza MS, Perri SH, Machado GF. Levels of matrix metalloproteinase-2 and metalloproteinase-9 in the cerebrospinal fluid of dogs with visceral leishmaniasis. *Parasite Immunol.* 2011;33:330-334.
230. Lugin J, Rosenblatt-Velin N, Parapanov R, Liaudet L. The role of oxidative stress during inflammatory processes. *Biol Chem.* 2014;395:203-230.

231. Heidarpour M, Soltani S, Mohri M, Khoshnegah J. Canine visceral leishmaniasis: relationships between oxidative stress, liver and kidney variables, trace elements, and clinical status. *Parasitol Res.* 2012;111:1491-1496.
232. Almeida BF, Narciso LG, Melo LM, et al. Leishmaniasis causes oxidative stress and alteration of oxidative metabolism and viability of neutrophils in dogs. *Vet J.* 2013;198:599-605.
233. Souza CC, Barreto Tde O, da Silva SM, et al. A potential link among antioxidant enzymes, histopathology and trace elements in canine visceral leishmaniasis. *Int J Exp Pathol.* 2014;95:260-270.
234. Paltrinieri S, Ravicini S, Rossi G, Roura X Serum concentration of the derivative of reactive oxygen metabolites (d-ROMs) in dogs with leishmaniosis. *Vet J.* 2010;186:393-395.
235. Oliva G, Roura X, Crotti A, et al. Guidelines for treatment of leishmaniasis in dogs. *J Am Vet Med Assoc.* 2010;236:1192-1199.
236. Bianciardi P, Brovida C, Valente M, et al. Administration of miltefosine and meglumine antimoniate in healthy dogs: clinicopathological evaluation of the impact on the kidneys. *Toxicol Pathol.* 2009;37:770-775.
237. Miró G, Oliva G, Cruz I, et al. Multicentric, controlled clinical study to evaluate effectiveness and safety of miltefosine and allopurinol for canine leishmaniosis. *Vet Dermatol.* 2009;20:397-404.
238. Xenoulis PG, Saridomichelakis MN, Chatzis MK, et al. Prospective evaluation of serum pancreatic lipase immunoreactivity and troponin I concentrations in *Leishmania infantum*-infected dogs treated with meglumine antimonate. *Vet Parasitol.* 2014;203:326-330.

239. Luciani A, Sconza S, Civitella C, Guglielmini C. Evaluation of the cardiac toxicity of N-methyl-glucamine antimoniate in dogs with naturally occurring leishmaniasis. *Vet J.* 2013;196:119-121.
240. Torres M, Bardagí M, Roura X, Zanna G, Ravera I, Ferrer L. Long term follow-up of dogs diagnosed with leishmaniosis (clinical stage II) and treated with meglumine antimoniate and allopurinol. *Vet J.* 2011;188:346-351.
241. Pierantozzi M, Roura X, Paltrinieri S, Poggi M, Zatelli A. Variation of proteinuria in dogs with leishmaniasis treated with meglumine antimoniate and allopurinol: 53 cases (2006-2010). *J Am Anim Hosp Assoc.* 2013;49:231-236.
242. Roura X, Fondati A, Lubas G, et al. Prognosis and monitoring of leishmaniasis in dogs: A working group report. *Vet J.* 2013;198:43-47.
243. Goldstein RE, Brovida C, Fernández-Del Palacio MJ, et al, Consensus recommendations for treatment for dogs with serology positive glomerular disease. *J Vet Intern Med.* 2013;27 Suppl 1:S60-6.
244. Rossi G, Ibba F, Meazzi S, Giordano A, Paltrinieri S. Paraoxonase activity as a tool for clinical monitoring of dogs treated for canine leishmaniasis. *Vet J.* 2014;199:143-149.
245. Rougier S, Hasseine L, Delaunay P, Michel G, Marty P. One-year clinical and parasitological follow-up of dogs treated with marbofloxacin for canine leishmaniosis. *Vet Parasitol.* 2012;186:245-253.
246. Manna L, Corso R, Galiero G, Cerrone A, Muzj P, Gravino AE. Long-term follow-up of dogs with leishmaniosis treated with meglumine antimoniate plus allopurinol versus miltefosine plus allopurinol. *Parasit Vectors.* 2015;8:289.
247. Morales-Yuste M, Morillas-Márquez F, Díaz-Sáez V, Barón-López S, Acedo-Sánchez C, Martín-Sánchez J. Epidemiological implications of the use of various methods for the diagnosis of canine leishmaniasis in dogs with different characteristics and in differing prevalence scenarios. *Parasitol Res.* 2012;111:155-164.